

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number  
WO 03/054228 A2(51) International Patent Classification<sup>7</sup>: C12Q 1/68

(74) Agents: GINZEL, Christian et al.: c/o Zimmermann &amp; Partner, Postfach 330 920, 80069 Munich (DE).

(21) International Application Number: PCT/EP02/14578

(22) International Filing Date:  
19 December 2002 (19.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/341,757 21 December 2001 (21.12.2001) US

(71) Applicant (for all designated States except US): AXXIMA PHARMACEUTICALS AG [DE/DE]; Am Klopferspitz 19, 82152 Martinsried (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SALASSIDIS, Konstadinos [GR/DE]; Hauptstrasse 14c, 85386 Eching (DE). SCHUBART, Daniel [DE/DE]; Avenue Charles de Gaulle 1c, 13469 Berlin (DE). GUTBROD, Heidrun [DE/DE]; Weberhofstrasse 8, 82166 Graefelfing (DE). MUELLER, Stefan [DE/DE]; Thalkirchnerstrasse 184, 81371 Munich (DE). KRAETZER, Friedrich [DE/DE]; Geisenbrunnerstrasse 51, 81475 Munich (DE). OBERT, Sabine [DE/DE]; Bellinzonastrasse 17, 81475 Munich (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HUMAN CELLULAR PROTEIN KINASES, METALLOPROTEASES AND PHOSPHATASES AS TARGETS FOR MEDICAL INTERVENTION AGAINST HEPATITIS C VIRUS INFECTIONS

(57) Abstract: The present invention relates to human cellular protein kinases, metalloproteases and one phosphatase: beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1 (NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649) as potential targets for medical intervention against Hepatitis C virus (HCV) infections. The present invention relates also to a method for the detection of compounds useful for prophylaxis and/or treatment of Hepatitis C virus infections, a method for detecting Hepatitis C virus infections in an individual or in cells. Mono- or polyclonal antibodies are disclosed effective for the treatment of HCV infections together with methods for treating Hepatitis C virus infections or for the regulation of Hepatitis C virus production and/or replication wherein said antibodies may be used. Finally the present invention relates to a solid support useful for detecting Hepatitis C virus infections or for screening compounds useful for prophylaxis and/or treatment of HCV infections.

WO 03/054228 A2

**Human cellular protein kinases, metalloproteases and phosphatases  
as targets for medical intervention against Hepatitis C virus infections**

5

**Specification**

The present invention relates to the human cellular protein kinases, metalloproteases and phosphatases:

- 10 1. beta-adrenergic receptor kinase 1 (NM\_001619, X61157, ADRBK1, GRK2, BARK1);
2. Mitogen activated protein kinase activated protein kinase 5 (AF032437, MAPKAPK5, PRAK, );
3. Insulin-stimulated protein kinase 1 (U08316, ribosomal protein S6 kinase 3  
15 90K);
4. Discoidin domain receptor family, member 1 (NM\_013994, DDR1, TRK E, NEP, CAK, X74979);
5. Protein Kinase C, mu (X75756, PKC-mu);
6. Protein Kinase C, theta (L01087, PKC-theta);
- 20 7. AMP-activated protein kinase beta 2 subunit (AJ 224538, AMPK beta 2);
8. JNK2 (U09759, L31951);
9. Human p21-activated protein kinase 2 (U24153, PAK2);
10. cyclin-dependent kinase 4 (U37022, cdk4);
11. MEK5 (U25265, Mitogen-activated protein kinase kinase 5);
- 25 12. MKP-L or DUSP14, formally also named as MKP-1 like tyrosine phosphatase (NM\_007026, AF038844);
13. ADAM22 (NM\_016351, AF155382; a disintegrin and metalloproteinase domain 22);
14. ADAM17 (U92649, XM\_002270; a disintegrin and metalloproteinase domain  
30 17);

as potential targets for medical intervention against Hepatitis C virus (HCV) infections. Furthermore, the present invention relates to a method for the detection of compounds useful for prophylaxis and/or treatment of Hepatitis C

virus infections and a method for detecting Hepatitis C virus infections in an individual or in cells. Also mono- or polyclonal antibodies are disclosed effective for the treatment of HCV infections together with methods for treating Hepatitis C virus infections or for the regulation of Hepatitis C virus production wherein said  
5 antibodies may be used. Finally the present invention relates to a solid support useful for detecting Hepatitis C virus infections or for screening compounds useful for prophylaxis and/or treatment of HCV infections.

## 10 **Background of the invention**

Hepatitis C Virus (HCV) infection is a major cause of chronic hepatitis, cirrhosis and hepatocellular carcinoma. The WHO estimates that approximately 3% of the world population, or 170 million people, have been infected with the Hepatitis C Virus. In the U.S., an estimated 3.9 million Americans have been infected (CDC  
15 fact sheet Sept. 00). Over 80% of HCV-infected individuals develop chronic hepatitis, which is associated with disease states ranging from asymptomatic carrier states to repeated inflammation of the liver and serious chronic liver disease. Over the course of 20 years, more than 20% of chronic HCV-patients are expected to be at risk to develop cirrhosis or progress to hepatocellular  
20 carcinoma. Liver failure from chronic hepatitis C is the leading indicator for liver transplantation. Excluding transplantation, the CDC estimates that medical and work-loss cost for HCV annually are around \$600 million. HCV is transmitted primarily by blood and blood products. Due to routine screening of the blood supplies from mid-1992, new transfusion-related cases are exceedingly rare and  
25 have been surpassed by injection drug use as the highest risk factor for acquiring the virus. There is also a sexual, however inefficient, route of transmission, and a 6% rate of transmission from infected mothers to their children, which is higher in case of HIV coinfection. In a certain percentage of infections, the mode of transmission remains unknown. In spite of the significant decline in incidence in  
30 the 1990's, the number of deaths (estimated deaths annually at the moment: 8000 to 10,000 in U.S.) and severe disease due to HCV is anticipated to triple in the next 10 to 20 years. (Sources: CDC fact sheet (accessed 12/12/00); Houghton M. Hepatitis C Viruses. In BN Fields, DM Knipe, PM Howley (ed.) Fields Virology.

1996. Lippencott-Raven Pub., Philadelphia; Rosen HR and Gretch DR, Molecular Medicine Today Vol5, 393, Sept. 99; Science 285, 26, July 99: News Focus: The scientific challenge of Hepatitis C; Wong JB et al, Am J Public Health, 90, 1562, Oct 2000: Estimating future hepatitis C morbidity, mortality, and costs in the United States).

According to the Consensus Statement from the EASL (European Association for the Study of the Liver) International Consensus Conference on Hepatitis C (February 26-28, 1999, Paris, France), combination therapy of alpha interferon and ribavirin is the recommended treatment for naive patients. Monotherapy with interferon has also been approved by the FDA, but the sustained response rate (HCV RNA remains undetectable in the serum for more than 6 months after end of therapy) is only 15 to 20%, in contrast to 35 to 45% with combination therapy. Interferons (Intron A, Schering-Plough; Roferon A, Hoffmann-LaRoche; Wellferon, Glaxo Wellcome; Infergen, Amgen) are injected subcutaneously three times a week, ribavirin (Rebetol, Schering-Plough) is an oral drug given twice a day. Recommended treatment duration is 6 to 12 months, depending on HCV genotype. Experimental forms of slow-release pegylated interferons (Pegasys, Hoffmann-LaRoche; PEG-Intron, Schering-Plough) have shown improvements in response rates (42 to 82% in combination with ribavirin) and application (once-weekly injection) in recent clinical studies (Hepatology32:4, Pt 2 of 2. Oct 2000; NEJM 343, 1673. Dec 00; NEJM 343, 1666. Dec 00). Common side effects of interferon therapy include: fatigue, muscle aches, head aches, nausea, fever, weight loss, irritability, depression, bone marrow suppression, reversible hair loss. The most common side effects of ribavirin are anemia, fatigue and irritability, itching, skin rash, nasal stuffiness, sinusitis, cough. More serious side effects of mono-and combination therapy occur in less than two percent of patients (NIDDK information: Chronic Hepatitis C: Current Disease Management. accessed 09.12.99). Some of the contraindications to interferon are psychosis or severe depression; neutropenia and/or thrombocytopenia; organ transplantation except liver; symptomatic heart disease; decompensated cirrhosis; uncontrolled seizures. Contraindications to ribavirin are end-stage renal failure; anemia;

hemoglobinopathies; severe heart disease; pregnancy; no reliable method of contraception (consensus statement EASL).

Experimental treatments that are not new forms of interferon are Maxamine  
5 (histamine dihydrochloride, Maxim Pharmaceuticals), which will be combined with  
Interferon in phase III studies, VX-497 (Vertex Pharmaceuticals), an IMP  
dehydrogenase inhibitor, as a less toxic ribavirin substitute in phase II, and  
amantadine (Endo Labs), an approved influenza drug, as the third component in  
triple therapy (phase II). Inhibitors for HCV enzymes such as protease inhibitors,  
10 RNA polymerase inhibitors, helicase inhibitors as well as ribozymes and antisense  
RNAs are under preclinical development (Boehringer Ingelheim, Ribozyme  
Pharmaceuticals, Vertex Pharmaceuticals, Schering-Plough, Hoffmann-LaRoche,  
Immusol, Merck etc.). No vaccine is available for prevention or therapeutic use,  
but several companies are trying to develop conventional or DNA vaccines or  
15 immunostimulatory agents (e.g. Chiron, Merck/Vical, Epimmune, NABI,  
Innogenetics).

In summary, the available treatment for chronic Hepatitis C is expensive, effective  
only in a certain percentage of patients and adverse side effects are not  
20 uncommon.

### **Description of the invention**

Recent research has revealed how cells communicate with each other to  
25 coordinate the growth and maintenance of the multitude of tissues within the  
human body. A key element of this communication network is the transmission of  
a signal from the exterior of a cell to its nucleus, which results in the activation or  
suppression of specific genes. This process is called signal transduction.

30 An integral part of signal transduction is the interaction of ligands, their receptors  
and intracellular signal transduction molecules. Ligands are messengers that bind  
to specific receptors on the surface of target cells. As a result of the binding, the  
receptors trigger the activation of a cascade of downstream signaling molecules,

thereby transmitting the message from the exterior of the cell to its nucleus. When the message reaches the nucleus, it initiates the modulation of specific genes, resulting in the production of RNA and finally proteins that carry out a specific biological function. Disturbed activity of signal transduction molecules  
5 may lead to the malfunctioning of cells and disease processes. Specifically, interaction of HCV with host cells is necessary for the virus to replicate.

The present invention is based upon the discovery of a group of human cellular protein kinases, metalloproteases and one phosphatase which are specifically up-  
10 or downregulated as a result of HCV replication in HCV infected host cells. The antiviral therapeutic research approach described herein focuses on discovering the cellular signal transduction pathways involved in viral infections. Identification of the signal transduction molecules, key to viral infection, provides for, among other things, novel diagnostic methods, for example, assays and compositions  
15 useful therefore, novel targets for antiviral therapeutics, a novel class of antiviral therapeutics, and new screening methods (e.g. assays) and materials to discover new antiviral agents.

This approach led to the development of a novel microarray platform technology,  
20 wherein a microarray of more than 900 signal transduction cDNAs was developed. This unique microarray technology was used to identify changes in RNA expression patterns (e.g. upregulation or downregulation) as a result of HCV infected host cells. Differential display techniques were used to pinpoint those signal transduction molecules useful as targets for drug intervention. Effective  
25 manipulation of these virally-controlled intracellular signal transduction pathways can alter (slow or stop altogether) the course of viral growth.

It is object of the present invention to provide novel targets for medical intervention, prophylaxis and/or treatment of Hepatitis C virus infections in  
30 mammals, including humans, and cells together with methods for detecting HCV infections in individuals and cells and methods for detecting compounds useful for prophylaxis and/or treatment of HCV infections. The object of the present

invention is solved by the teaching of the independent claims. Further preferred embodiments are disclosed in the dependent claims.

It is now revealed for the first time that the human cellular proteins

- 5 beta-adrenergic receptor kinase 1 (gene accession number NM\_001619, other names: X61157, ADRBK1, GRK2, BARK1),  
Mitogen activated protein kinase activated protein kinase 5 (also known as: AF032437, MAPKAPK5, or PRAK),  
Insulin-stimulated protein kinase 1 (also known as: U08316, ribosomal protein S6  
10 kinase 3 90K),  
Discoidin domain receptor family, member 1 (gene accession number NM\_013994, other names: X74979, DDR1, TRK E, NEP, or CAK),  
Protein Kinase C, mu (also known as: X75756 or PKC-mu),  
Protein Kinase C, theta (also known as: L01087 or PKC-theta),  
15 AMP-activated protein kinase beta 2 subunit (also known as: AJ 224538 or AMPK beta 2),  
JNK2 (also known as: U09759 or L31951),  
Human p21-activated protein kinase 2 (also known as: U24153 or PAK2),  
cyclin-dependent kinase 4 (also known as: U37022 or cdk4),  
20 MEK5 (also known as: U25265 or Mitogen-activated protein kinase kinase 5),  
MKP-L (gene accession number: NM\_007026; also known as AF038844),  
ADAM22 (gene accession number: NM\_016351, a disintegrin and metalloproteinase domain 22, also known as AF155382),  
ADAM17 (also known as: U92649 or a disintegrin and metalloproteinase domain  
25 17 (tumor necrosis factor, alpha, converting enzyme) also known as XM\_002270),  
are specifically and uniquely up- or downregulated in a cell as a result of HCV infection. These cellular protein kinases, metalloproteases and the phosphatase therefore identify novel diagnostic and therapeutic targets for HCV infection.
- 30 The only reliable experimental HCV infection studies have been performed with chimpanzees. There is no simple cell culture infection system available for HCV. Although a number of reports have been published describing *in vitro* propagation attempts of HCV in primary cells and cell lines, questions remain concerning

reproducibility, low levels of expression and properly controlled detection methods (reviewed in J. Gen Virol. 81, 1631; Antiviral Chemistry and Chemotherapy 10, 99). Thus, the replicon system described by Bartenschlager and coworkers (Lohmann et al, Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. Science 285, 110. 1999) was used for the studies disclosed herein. This replicon system reproduces a crucial part of the HCV replication cycle which is used as a system for simulating HCV infection. Bartenschlager's group produced bicistronic recombinant RNAs, so-called "replicons", which carry the Neomycin-phosphotransferase gene as well as a version of the HCV genome where the sequences for the structural HCV proteins were deleted. After transfection of the subgenomic HCV RNA molecules into the human hepatoma cell line Huh-7, cells supporting efficient RNA-dependent RNA replication of the HCV replicons were selected based on co-amplification of the neo gene and resulting resistance to the antibiotic G418. Integration of coding information into the cellular genome was an exclusion criterium for functional replicons. Several lines were established from G418 resistant clones with autonomously replicating HCV RNAs detectable by Northern blot. Minus-strand RNA replication intermediates were detected by Northern blot or metabolic radio-labeling, and the production of nonstructural HCV proteins was demonstrated by immuno-precipitation after metabolic labeling or Western blot.

Possible influences and/or dependencies of HCV's RNA-dependent RNA replication and nonstructural proteins on host cell transcription are accessible to analysis with the cDNA arrays used in the inventive methods described herein. Expression levels can be confirmed using Northern or Taqman analysis at the RNA and Western blot analysis at the protein level. Huh-pcDNA3 cells are Huh7 cells resistant to G418 by integration of a plasmid and serve as negative control. Three replicon lines were analyzed for changes in cellular RNA expression patterns compared to the control line:

- Huh-9-13: cell line with persistent replicon I377/NS3-3'/wt, described in Science 1999, 285, 110-113,
- Huh-5-15: cell line with persistent replicon I389/NS3-3'/wt, described in Science 1999, 285, 110-113,
- Huh-11-7: cell line with persistent replicon I377/NS2-3'/wt, described in Science 1999, 285, 110-113.



Based on the discoveries reported herein, one aspect of the present invention is directed to a screening method for detecting compounds useful for the prophylaxis and/or treatment of Hepatitis C virus infections. Specifically, this method involves

5 contacting a test compound with at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein

10 Kinase C, mu (X75756), Protein Kinase C, theta(L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (AF038844, NM\_007026), ADAM22 (NM\_016351), ADAM17 (U92649) and detecting the human cellular protein kinase, metalloprotease or phosphatase

15 activity.

Another aspect of the present invention is directed to a diagnostic method, an assay for detecting Hepatitis C virus infections in an individual or cells. This method involves providing a sample from the individual or providing cells and detecting

20 activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of: beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein

25 Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), ADAM17 (U92649) and detecting the human cellular protein kinase, metalloprotease or phosphatase activity.

30

Accordingly, one aspect of the present invention is directed to novel compounds useful in the above-identified methods. Therefore, the present invention relates to a monoclonal or polyclonal antibody that binds to a human cellular protein kinase,

metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein  
5 Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351) and ADAM17 (U92649).

10 Furthermore, the present invention discloses a method for treating Hepatitis C virus infection in an individual comprising the step of administering a pharmaceutically effective amount of an agent which inhibits at least partially the activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from  
15 the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4  
20 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

Another object of the present invention is to provide a method for regulating the production of Hepatitis C virus in cells comprising the step of administering a  
25 pharmaceutically effective amount of an agent to said cells wherein said agent inhibits at least partially the activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1  
30 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L

( NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649). The above-mentioned monoclonal or polyclonal antibodies directed against these targets may be used as pharmaceutically active agents within said methods.

- 5 In order to identify HCV infections and new inhibitors and new pharmaceutically active compounds against Hepatitis C viruses a further aspect of the present invention is directed to a solid support useful for detecting Hepatitis C virus infections in an individual or in cells comprising an immobilized oligonucleotide, wherein said oligonucleotide is capable of detecting activity of at least one human
- 10 cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated
- 15 protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649). Said solid support is also useful to screen compounds for the prophylaxis and/or treatment of Hepatitis C virus infections in an individual comprising at least one immobilized
- 20 oligonucleotide, wherein said oligonucleotide encodes one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein
- 25 Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649) or comprising at least one immobilized human cellular protein kinase, metalloprotease or
- 30 phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein

Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

5

Yet another aspect of the present invention is directed to a novel therapeutic composition useful for the prophylaxis and/or treatment of an individual afflicted with Hepatitis C virus comprising at least one agent capable of inhibiting activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected  
10 from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759),  
15 Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

## 20 Detailed description of the invention

Utilizing microarray technology, a unique microarray of more than 900 signal transduction cDNAs was developed. This array was used to compare signal transduction mRNA expression patterns (e.g. upregulation or downregulation) from HCV Replicon cells Huh-9-13, Huh-5-15, and Huh-11-7 with Huh-pcDNA  
25 control cells which do not contain HCV Replicons. These HCV Replicon cells serve as a system for simulation of HCV infected cell systems, especially for simulating HCV infected mammals, including humans. Interference of HCV with the cellular signaling events is reflected in differential gene expression when compared to cellular signaling in control cells. Results from this novel signal  
30 transduction microarray analysis revealed significant up- or downregulation of human cellular protein kinases, metalloproteases and one phosphatase. Radioactively labeled complex cDNA-probes from HCV Replicon cells Huh-9-13, Huh-5-15, and Huh-11-7 were hybridized to cDNA-arrays and compared to

hybridizations with cDNA-probes from Huh-pcDNA control cells which did not contain HCV Replicons. Surprisingly it was found that the following cellular targets are significantly up- or downregulated:

	beta-adrenergic receptor kinase 1 (NM_001619):	2.7 – 3.5 fold stronger
5	Mitogen activated protein kinase activated protein kinase 5 (AF032437):	2.2 – 3.0 fold stronger
	Insulin-stimulated protein kinase 1 (U08316):	2.2 – 3.1 fold stronger
	TRK E (NM_013994):	3.2 -10.3 fold stronger
	Human p21-activated protein kinase 2 (U24153):	1.8 – 2.7 fold stronger
10	PKC-mu (X75756):	2.3 – 3.2 fold weaker
	PKC-theta (L01087):	2.6 – 3.3 fold weaker
	AMP-activated protein kinase beta 2 subunit (AJ 224538):	1.9 – 2.2 fold weaker
	JNK2 (U09759):	2.6 – 4.1 fold weaker
15	cdk4 (U37022)	1.8 – 3.3 fold stronger
	MEK5 (U25265)	0.9 – 3.6 fold stronger
	MKP-L (NM_007026):	2.1 – 2.3 fold weaker
	ADAM22 (NM_016351):	2.5 – 3.8 fold weaker
20	ADAM17 (U92649):	3.4 – 3.8 fold weaker

Disclosed herein is the first report describing the role of human cellular proteins beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1 (NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein

kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), ADAM17 (U92649) in the signal transduction of the HCV infection process. As a result of these discoveries, novel compounds and inhibitors against the above-mentioned human cellular protein kinases, metalloproteases and phosphatases may be found by the use of the inventive methods disclosed herein.

ADAM17 and ADAM22 are proteins of the ADAM family (proteins containing a disintegrin and metalloprotease domain). ADAM17 is also known as "Homo sapiens a disintegrin and metalloproteinase domain 17" and ADAM22 is known as "Homo sapiens metalloprotease-like, disintegrin-like, cysteine-rich protein 2 epsilon".

As used herein, the term "inhibitor" refers to any compound capable of downregulating, decreasing, suppressing or otherwise regulating the amount and/or activity of the human cellular proteins beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1 (NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649). Generally, human cellular protein inhibitors may be proteins, oligo- and polypeptides, nucleic acids, small chemical molecules, or other chemical moieties.

As used herein, the term "regulating expression and/or activity" generally refers to any process that functions to control or modulate the quantity or activity (functionality) of a cellular component. Static regulation maintains expression and/or activity at some given level. Upregulation refers to a relative increase in expression and/or activity. Accordingly downregulation refers to a relative decrease in expression and/or activity. In the present invention, regulation is preferably downregulation of a cellular component. As used herein,

downregulation is synonymous with inhibition of a given cellular component's activity.

Therapeutics, pharmaceutically active agents or inhibitors, respectively, may be administered to cells from an individual *in vitro*, or may involve *in vivo* administration to the individual. The term "individual" preferably refers to mammals and most preferably to humans. Routes of administration of pharmaceutical preparations to an individual may include oral and parenteral, including dermal, intradermal, intragastral, intracutan, intravasal, intravenous, intramuscular, intraperitoneal, intranasal, intravaginal, intrabuccal, percutan, rectal, subcutaneous, sublingual, topical or transdermal application, but are not limited the these ways of administration. For instance, the preferred preparations are in administratable form which is suitable for oral application. These administratable forms, for example, include pills, tablets, film tablets, coated tablets, capsules, powders and deposits. Administration to an individual may be in a single dose or in repeated administrations, and may be in any of a variety of physiologically acceptable salt forms, and/or with an acceptable pharmaceutical carrier, binder, lubricant, excipient, diluent and/or adjuvant. Pharmaceutically acceptable salt forms and standard pharmaceutical formulation techniques are well known to persons skilled in the art.

As used herein, a "pharmaceutical effective amount" of a human cellular protein kinase, metalloprotease or phosphatase inhibitor is an amount effective to achieve the desired physiological result, either in cells treated *in vitro* or in a subject treated *in vivo*. Specifically, a pharmaceutically effective amount is an amount sufficient to inhibit, for some period of time, one or more of the clinically defined pathological processes associated with the viral infection. The effective amount may vary depending on the specific human cellular protein kinase, metalloprotease or phosphatase inhibitor selected, and is also dependent on a variety of factors and conditions related to the subject to be treated and the severity of the infection. For example, if the inhibitor is to be administered *in vivo*, factors such as the age, weight and health of the patient as well as dose response curves and toxicity data obtained in pre-clinical animal work would be among

those considered. If the inhibitor is to be contacted with the cells *in vitro*, one would also design a variety of pre-clinical *in vitro* studies to assess such parameters as uptake, half-life, dose, toxicity, etc. The determination of a pharmaceutically effective amount for a given agent is well within the ability of  
5 those skilled in the art.

It is also apparent to a person skilled in the art that detection includes any method known in the art useful to indicate the presence, absence, or amount of a detection target. Such methods may include, but are not limited to, any molecular  
10 or cellular techniques, used singularly or in combination, including, but not limited to: hybridization and/or binding techniques, including blotting techniques and immunoassays; labeling techniques (chemiluminescent, colorimetric, fluorescent, radioisotopic); spectroscopic techniques; separations technology, including precipitations, electrophoresis, chromatography, centrifugation, ultrafiltration, cell  
15 sorting; and enzymatic manipulations (e.g., digestion).

The present disclosure teaches for the first time the up- or downregulation of a group of human cellular protein kinases, metalloproteases and a phosphatase specifically involved in the viral infection of Hepatitis C virus. Thus, the present  
20 invention is also directed to a method useful for detecting novel compounds useful for prophylaxis and/or treatment of HCV infections.

Methods of the present invention identify compounds useful for prophylaxis and/or treatment of HCV infections by screening a test compound, or a library of test  
25 compounds, for its ability to inhibit any one or more of the group of human cellular protein kinases, metalloproteases or phosphatases identified herein as characteristically up- or downregulated during HCV growth and RNA replication inside a cell. A variety of assay protocols and detection techniques are well known in the art and easily adapted for this purpose by a skilled practitioner. Such  
30 methods include, but are not limited to, high throughput assays (e.g., microarray technology, phage display technology), and *in vitro* and *in vivo* cellular and tissue assays.



In a related aspect, the present invention provides, in view of the discovery of human cellular protein kinases, metalloproteases and phosphatases specifically involved in the HCV infection process, an assay component specially useful for detecting HCV in an individual or in cells. Preferably the assay component  
5 comprises oligonucleotides immobilized on a solid support capable of detecting activity of one or more of the human cellular protein kinases, metalloproteases or phosphatase comprising:

the kinases	beta-adrenergic receptor kinase 1 (NM_001619), Mitogen activated protein kinase activated protein kinase 5 10 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1 (NM_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21- 15 activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265);
the metalloproteases	ADAM22 (NM_016351), ADAM17 (U92649) and
the phosphatase	MKP-L (NM_007026).

20 Preferably the solid support would contain oligonucleotides of sufficient quality and quantity to detect all of the above-mentioned human cellular proteins (e.g., a nucleic acid microarray).

Similarly, it is an object of the present invention to provide an assay component  
25 specially useful for screening compounds useful for the prophylaxis and/or treatment of HCV infections. One preferred assay component comprises oligonucleotides that encode one or more human cellular protein kinases: beta-adrenergic receptor kinase (X61157), Mitogen activated protein kinase activated  
protein kinase (AF032437), Insulin-stimulated protein kinase 1 (U08316), TRK E  
30 (X74979), PKC-mu (X75756), PKC-theta (L01087), AMP-activated protein kinase  
beta 2 subunit (AJ 224538), JNK2 (L31951), Human p21-activated protein kinase,  
and PAK2 (U24153); or metalloproteases: ADAM22 (AF155382) and ADAM17

(XM\_002270); or the Phosphatase MKP-L, (NM\_007026) immobilized on a solid support.

5 In another embodiment, the assay component comprises peptide fragments of one or more of the above-identified human cellular proteins immobilized on a solid support. Once again the most preferred solid support embodiment would contain polymers of sufficient quality and quantity to detect all of the above-mentioned human cellular protein kinases, metalloproteases and the phosphatase (e.g., a nucleic acid or a peptide microarray). A variety of supports and constructions of  
10 the same for the methods disclosed herein are well known in the art and easily adapted for this purpose by a skilled practitioner (cf., for example: Marschall, 1999 "Do-it-yourself gene watching" Science 286, 444-447; Service 2000 "Protein arrays step out of DNA's shadow" Science 289, 1673).

15 It is preferred that mRNA is measured as an indication of expression. Methods for assaying for mRNA include, but are not limited to, Northern blots, slot blots, dot blots, and hybridization to an ordered array of oligonucleotides. Nucleic acid probes useful for assay of a sample are preferably of sufficient length to specifically hybridize only to appropriate, complementary transcripts. Typically the  
20 oligonucleotide probes will be at least 10 to 25 nucleotides in length. In some cases longer probes of at least 30, 40, or 50 up to 2500 nucleotides will be desirable.

The polypeptide product of gene expression may be assayed to determine the  
25 amount of expression as well. Methods for assaying for a protein include, but are not limited to, western blot, immuno-precipitation, radioimmuno assay, and peptide immobilization in an ordered array. It is understood, however, that any method for specifically and quantitatively measuring a specific protein or mRNA product can be used.

30

A variety of supports upon which nucleic acids or peptides can be immobilized are known in the art, for example filters, or polyvinyl chloride dishes. Any solid surface to which oligonucleotides or peptides can be bound, either directly or indirectly,

either covalently or non-covalently, can be used. A preferred solid support is a microarray membrane filter or a "biochip". These contain particular polymer probes in predetermined locations on the array. Each predetermined location may contain more than one molecule of the probe, but each molecule within the  
5 predetermined location has an identical sequence.

The present invention incorporates by reference in their entirety techniques well known in the field of molecular biology. These techniques include, but are not limited to, techniques described in the following publications:

- 10 Ausubel, F.M. et al. eds., "Short Protocols In Molecular Biology" 4<sup>th</sup> Ed. 1999, John Wiley & Sons, NY (ISBN 0-471-32938-X);
- Old, R.W. & S.B. Primrose "Principles of Gene Manipulation: An Introduction To Genetic Engineering" 3<sup>rd</sup> Ed. 1985, Blackwell Scientific Publications, Boston. Studies in Microbiology: V.2, 409 pp. (ISBN 0-632-01318-4);
- 15 Miller, J.H. & M.P. Calos eds., "Gene Transfer Vectors For Mammalian Cells" 1987, Cold Spring Harbor Laboratory Press, NY. 169 pp. (ISBN 0-87969-198-0);
- Mayer, R.J. & J.H. Walker eds. "Immunochemical Methods In Cell and Molecular Biology" 1987, Academic Press, London. 325 pp. (ISBN 0-12480-855-7);
- Sambrook, J. et al. eds., "Molecular Cloning: A Laboratory Manual" 2<sup>nd</sup> Ed. 1989,  
20 Cold Spring Harbor Laboratory Press, NY. Vols. 1-3. (ISBN 0-87969-309-6);
- Winnacker, E.L. "From Genes To Clones: Introduction To Gene Technology" 1987 VCH Publishers, NY. (translated by Horst Ibelgaufts) 634 pp. (ISBN 0-89573-614-4).

25 The present invention further incorporates by reference in their entirety techniques well known in the field of microarray construction and analysis. These techniques include, but are not limited to, techniques described in the following patents and patent applications describing array of biopolymeric compounds and methods for their fabrication:

- 30 U.S. Pat. Nos. 5,242,974; 5,384,261; 5,405,783; 5,412,087; 5,424,186;  
5,429,807; 5,436,327; 5,445,934; 5,472,672; 5,527,681; 5,529,756;  
5,545,531; 5,554,501; 5,556,752; 5,561,071; 5,559,895; 5,624,711;

5,639,603; 5,658,734; 5,807,522; 6,087,102; WO 93/17126; WO 95/11995; WO 95/35505; EP 742 287; and EP 799 897.

Techniques also include, but are not limited to, techniques described in the following patents and patent application describing methods of using arrays in various applications:

U.S. Pat. Nos. 5,143,854; 5,288,644; 5,324,633; 5,432,049;  
5,470,710; 5,492,806; 5,503,980; 5,510,270; 5,525,464; 5,547,839;  
5,580,732; 5,661,028; 5,994,076; 6,033,860; 6,040,138; 6,040,140;  
10 WO 95/21265; WO 96/31622; WO 97/10365; WO 97/27317;  
EP 373 203; and EP 785 280

It is readily apparent to those skilled in the art that other suitable modifications and adaptations of the compositions and methods of the invention described herein are evident and may be made without departing from the scope of the invention or the embodiments disclosed herein. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included for purposes of illustration only and are not intended to be limiting of the invention.

20

## Examples

### Materials and Methods

25

#### 1. Generation of cDNA-arrays on membranes

In order to manufacture cDNAs-arrays on membranes, the following strategy was pursued: cDNAs encoding parts of or full length proteins of interest – in the following referred to as “target cDNAs” – were cloned into the plasmid Bluescript II KS<sup>+</sup> (Stratagene, USA). Large scale purifications of these plasmids were performed according to standard techniques and 200 µl aliquots (1 µg/µl plasmid concentration) were transferred into appropriate 96 well plates. Plates were closed with sealing tape and chilled on ice for 5 minutes after incubation for 10 minutes at 95°C. 10 µl of 0.6N NaOH were added and the mix was stored for 20

30

minutes at room temperature before addition of 10  $\mu$ l 2.5M Tris-HCl pH 7.1 and 20  $\mu$ l 40x SSC. Target cDNAs were spotted onto Nylon or Nitrocellulose membranes using a BioGrid (BioRobotics, UK) equipped with a 0.7 mm pintool. In this way, between 200 ng and 350 ng of plasmids encoding target cDNAs were transferred  
5 onto the membranes and crosslinked to the membranes by ultraviolet light ( $1.2 \times 10^5 \mu\text{J}/\text{cm}^2$ ). The arrays were stored for use in subsequent experiments at at 4°C.

## 2. Cellular HCV RNA replication system

10 Huh-pcDNA3, Huh-9-13, Huh-5-15 and Huh-11-7 cells were grown in DMEM supplemented with 10% FCS, 2 mM Glutamine, Penicillin (100 IU/ml) / Streptomycin (100  $\mu$ g/ml) and 1x nonessential amino acids in the presence of 1 mg/ml G418. Cells were routinely passaged three times a week at a dilution of 1:3 or 1:2.

## 3. Lysis of cells, and isolation of total RNA

15 Huh-pcDNA3, Huh-9-13, Huh-5-15 and Huh-11-7 cells were seeded at  $5 \times 10^5$  cells per 10 cm plate in medium without G418. The medium was changed 3 days after plating and cells were harvested 5 days after plating by lysing the cells  
20 directly on the plate with 4 ml of Tri reagent (Molecular Research Center, Inc., USA). The lysates were stored at room temperature for 5 minutes and then centrifuged at 12000xg for 15 minutes at 4°C. The supernatant was mixed with 0,1 ml of 1-Bromo-3-chloropropane per 1 ml of Tri reagent and vigorously shaken. The suspension was stored for 5 minutes at room temperature and then  
25 centrifuged at 12000xg for 15 minutes at 4°C. The colorless upper phase was transferred into new tubes, mixed with 5  $\mu$ l of polyacryl-carrier (Molecular Research Center Inc., USA) and with 0.5 ml of isopropanol per 1 ml of Tri reagent and vigorously shaken. The samples were stored at room temperature for 5 minutes and then centrifuged at 12000xg for 8 minutes at 4°C. The supernatant  
30 was removed and the RNA pellet washed twice with 1 ml of 75% Ethanol. The pellet was dried and resuspended in 25  $\mu$ l of RNase-free buffer per initial 1 ml lysate.

#### 4. Preparation of radioactively labelled cDNA probes from RNA

In order to obtain a radioactively labeled cDNA probe, RNA was transcribed into a cDNA-probe in the presence of radioactively labeled dATP. 12 µl bidistilled DEPC treated H<sub>2</sub>O containing 1 µg of primer TXN (5'-TTT TTT TTT TTT TTT TXN-3' with T → dTTP; N → dATP, dCTP, dGTP or dTTP; X → dATP, dCTP or dGTP) and total RNA (6 µg) were shaken between 5 and 15' at 60°C and then incubated at 4°C for 2 to 10 minutes. After centrifugation (30 seconds, 10000xg) 7 µl of a mix consisting of 100 µCi dATP-P<sup>33</sup> (Amersham, UK) which were dried under vacuum previously and resuspended in 4 µl first strand buffer (Life Technologies, USA), 2 µl 0.1M DTT and 1 µl labeling solution (4mM dCTP, dGTP, dTTP each and 80 µM dATP final concentration) were added. Following the addition of 1 µl Superscript II reverse transcriptase (Life Technologies, USA) the reaction was incubated for 10 minutes at room temperature and then for 60 minutes at 38°C. Subsequently, the reaction was vigorously shaken for 30 minutes at 68°C after adding 5 µl 0.5M EDTA and 25 µl 0.6M NaOH.

Unincorporated nucleotides were removed from the labeling reaction using ProbeQuant G-50 columns (Amersham, UK). The column was vigorously shaken and centrifuged for 1 minute at 735xg in an appropriate reaction tube after bottom closure and lid were removed. The column was placed into a new reaction tube, the probe was applied onto the center of the column material and the column was centrifuged for 2 minutes at 735xg. The flow-trough was transferred into new reaction tubes and filled up to a volume of 100 µl with 10mM Tris, pH 7.4, 1 mM EDTA. The probe was precipitated by centrifugation for 15 minutes at 12000xg after 4 µl of 5M NaCl, 1 µl polyacryl-carrier (Molecular Research Center Inc., USA) and 250µl Ethanol were added. The supernatant was discarded and the pellet dried before starting with the hybridisation.

30

#### 5. Hybridisation of radioactively labeled cDNA-probes to cDNA-arrays

The pellet was resuspended in 10 µl C<sub>0</sub>T DNA (1 µg/µl, Roche Diagnostics, Germany), 10 µl yeast tRNA (1 µg/µl Sigma, USA) and 10 µl polyA (1 µg/µl, Roche

Diagnostics, Germany). Herring sperm DNA was added to a final concentration of 100 µg/ml and the volume was filled up to 100 µl with 5 µl 10% SDS (Sodiumdodecylsulfat), 25 µl 20x SSPE and bidistilled H<sub>2</sub>O. The mix was put on 95°C for 5 minutes, centrifuged for 30 seconds at 10000xg and vigorously shaken for 60 minutes at 65°C. A 1 µl aliquot of the probe was used to measure the incorporation of radioactive dATP with a scintillation counter. Probes with at least a total of 20x10<sup>6</sup> cpm were used. The arrays were prehybridised for at least 3 hours at 65°C in hybridisation solution in a roller bottle oven. After prehybridisation the radioactively labeled probe was added into the hybridisation solution and hybridisation was continued for 20 hours. The probe was discarded and replaced with wash solution A (2xSSC). The arrays were washed twice in wash solution A at room temperature in the roller oven. Afterwards, wash solution A was replaced by wash solution B (2x SSC, 0.5% SDS) preheated to 65°C and arrays were washed twice for 30 minutes at 65°C. Then, wash solution B was replaced by wash solution C (0.5x SSC, 0.5% SDS) preheated to 65°C and arrays were washed twice for 30 minutes at 65°C. The moist arrays were wrapped in airtight bags and exposed for 8 to 72 hours on erased phosphorimager screens (Fujifilm, Japan).

## 20 6. Analysis of cDNA-arrays

The exposed phosphorimager screens were scanned with a resolution of 100 µ and 16bits per pixel using a BAS-1800 (Fujifilm, Japan). Files were imported into the computer program ArrayVision (Imaging Research, Canada). Using the program's features, the hybridisation signals of each target cDNA were converted into numbers. The strength of the hybridisation signals reflected the quantity of RNA molecules present in the probe. Differentially expressed genes were selected according to the ratio of their signal strength after normalization to the overall intensity of the arrays.

30

## 7. Analysis of expression levels by Northern blot experiment

Huh-pcDNA3, Huh-9-13, Huh-5-15 and Huh-11-7 cells were seeded at 5 x 10<sup>5</sup> cells per 10 cm plate in medium without G148. Cells were harvested after 3 days

by lysing the cells directly on the plate with 4 ml of Tri reagent (Molecular Research Centre, Inc., USA). The lysates were stored at room temperature for 5 minutes and then centrifuged at 12000xg for 15 minutes at 4°C. The supernatant was mixed with 0,1 ml of 1-bromo-3-chloropropane per 1 ml of Tri reagent and vigorously shaken. The suspension was stored for 5 minutes at room temperature and then centrifuged at 12000xg for 15 minutes at 4°C. The colourless upper phase was transferred into new tubes, mixed with 5 µl of poly-acryl-carrier (Molecular Research Centre, Inc., USA) and with 0.5 ml of isopropanol per 1 ml of Tri reagent and vigorously shaken. The samples were stored at room temperature for 5 minutes and then centrifuged at 12000xg for 8 minutes at 4°C. The supernatant was removed and the RNA pellet washed twice with 1 ml of 75% Ethanol. The pellet was dried and resuspended in 25 µl of RNase-free water per initial 1 ml lysate. 8 µg of total RNA per sample was loaded onto formaldehyde-containing agarose gels (Sambrook et al. Cloning manual, CSHL press, 1989) and transferred to HYbond NX membranes (Amersham) overnight in 20x SSC (3M NaCl, 300mM C<sub>6</sub>H<sub>5</sub>Na<sub>3</sub>O<sub>7</sub> x 2 H<sub>2</sub>O, pH 7.0) by capillary transfer. RNA was immobilized to the filter using UV-crosslinking (120mJ/cm<sup>2</sup> for 25 seconds). Filters were hybridized to oligonucleotide probes or random-primed probes specific for the genes in question. Quantitation of signals was performed with a Fuji phosphoimager.

#### 8. Verification of de-regulated genes by quantitative Real-Time PCR

Quantitative RT-PCR was used to verify hits resulting from DNA macroarray experiments by exploiting the 5'-exonuclease of Taq DNA polymerases to cleave the 5' fluorescent label of an oligonucleotide. Total RNA was extracted from cell lines (Qiagen RNeasy Mini Kit, QIAGEN, Hilden) and was reverse transcribed with Superscript II (Invitrogen, Karlsruhe) according to the manufacturer's protocol with 5 µg of RNA as a template and oligodT primers. Subsequently, the cDNA was analysed on a ABI PRISM 7000 Sequence Detection System (Applied Biosystems, Darmstadt) with the 5'exonuclease assay by using the TaqMan Universal PCR Master Mix (#4324018, Applied Biosystems, Darmstadt) and non-extendible oligonucleotides. Gene-specific TaqMan probes were labelled with the reporter dye FAM<sup>TM</sup> at the 5'-end and the quencher dye TAMRA<sup>TM</sup> at the 3' end of



the probe. GAPDH and 18SrRNA were used as reference genes with TaqMan probes that were labelled with VIC<sup>TM</sup> and TAMRA<sup>TM</sup> accordingly.

Experimental conditions were 2 minutes 50°C, 10 minutes 95°C, followed by 40 cycles with 15 seconds at 95°C and 1 minute at 60°C. Primer Express software  
5 was used to design primers with a melting temperature of 58 - 60°C amplifying an amplicon of a maximum length of 150 bp.

#### 9. Analysis of expression levels by Western blot experiments

Huh-pcDNA3, Huh-9-13, Huh-5-15 and Huh-11-7 cells were seeded at  $5 \times 10^5$   
10 cells per 10 cm plate in medium without G148. Cells were harvested after 3 days by the addition of 500 µl of 1x SDS sample buffer (62.5 mM Tris-HCl pH 6.8, 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% bromphenol blue) or RIPA lysis buffer. Lysates were separated on SDS-poly acrylamide gels and proteins transferred to nitrocellulose. Western blotting was performed with the appropriate  
15 antibodies according to the manufacturers instructions.

### Claims

1. A method for detecting compounds useful for the prophylaxis and/or  
5 treatment of Hepatitis C virus infections, the method comprising the following steps:
- a) contacting a test compound with at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619),  
10 Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), ADAM17 (U92649); and
- 15 b) determining the activity of said human cellular protein kinase, metalloprotease or phosphatase.
- 20
2. A method for detecting Hepatitis C virus infections in an individual, the method comprising the following steps:
- a) providing a sample from said individual; and
- b) determining the activity in said sample of at least one human cellular  
25 protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4  
30

(U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

3. A method for detecting Hepatitis C virus infections in cells, the method  
5 comprising the following steps:
  - a) providing said cells; and
  - b) determining the activity in said cells of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).  
10  
15
4. An antibody that binds to a human cellular protein kinase, metalloprotease or  
20 phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).  
25
5. The antibody of claim 4, wherein said antibody is a monoclonal or polyclonal  
30 antibody.
6. A method for treating Hepatitis C virus infection in an individual, the method comprising the step of administering a pharmaceutically effective amount of an

agent which inhibits at least partially the activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

7. A method for regulating the production of Hepatitis C virus in cells, the method comprising the step of administering a pharmaceutically effective amount of an agent to said cells wherein said agent inhibits at least partially the activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

8. The method according to claim 6 or 7, wherein the agent is an antibody which binds to at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein

kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).

- 5 9. The method according to claim 8, wherein said antibody is a monoclonal or polyclonal antibody.
- 10 10. A solid support useful for detecting Hepatitis C virus infections in an individual, the solid support comprising an immobilized oligonucleotide, wherein said oligonucleotide is capable of detecting activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).
- 20 11. A solid support useful for detecting Hepatitis C virus infections in cells, the solid support comprising an immobilized oligonucleotide, wherein said oligonucleotide is capable of detecting activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).
- 25 30

12. A solid support useful for screening compounds useful for the prophylaxis and/or treatment of Hepatitis C virus infections in an individual, the solid support comprising at least one immobilized oligonucleotide, wherein said oligonucleotide encodes one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).
13. A solid support useful for screening compounds useful for the prophylaxis and/or treatment of Hepatitis C virus infections in an individual, the solid support comprising at least one immobilized human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and ADAM17 (U92649).
14. A composition useful for the prophylaxis and/or treatment of an individual afflicted with Hepatitis C virus, the composition comprising at least one agent capable of inhibiting activity of at least one human cellular protein kinase, metalloprotease or phosphatase selected from the group consisting of beta-adrenergic receptor kinase 1 (NM\_001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated protein kinase 1 (U08316), Discoidin domain receptor family, member 1(NM\_013994), Protein

Kinase C, mu (X75756), Protein Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM\_007026), ADAM22 (NM\_016351), and  
5 ADAM17 (U92649).

# 10. Nucleic acid and amino acid sequences of human cellular proteins:

DEFINITION Homo sapiens adrenergic, beta, receptor kinase 1 (ADRBK1), mRNA.

5 ACCESSION NM\_001619  
EC number = "2.7.1.126"

10 DEFINITION Homo sapiens mitogen activated protein kinase activated  
protein  
kinase gene, complete cds.  
15 ACCESSION AF032437  
VERSION AF032437.1 GI:3133290  
FEATURES Location/Qualifiers  
source 1..1968  
/organism="Homo sapiens"  
/db\_xref="taxon:9606"  
20 CDS 192..1607  
/note="MAPK-activated protein kinase; PRK"  
/codon\_start=1  
/product="mitogen activated protein kinase activated  
protein kinase"  
25 /protein\_id="AAC39863.1"  
/db\_xref="GI:3133291"

/translation="MSEESDMDKAIKETSILEEYSINWTQKLGAGISGPVRVCVKKST

30 QERFALKILLDRPKARNEVRLHMMCATHPNIVQIIEVFANSVQFPHESSPRARLLIVM  
EMMEGGELFHRISQHRHFTEKQASQVTKQIALALRHCHLLNIAHRDLKPENLLFKDNS  
LDAPVKLCDFGFAKIDQGLMTPQFTPPYVAPQVLEAQRHQKEKSGIIPTSPTPTY  
35 NKSCDLWSLGVIIYVMLCGYPPFYSKHHSRTIPKDMRRKIMTGSFEFPEEWSQISEM  
AKDVVRKLLKVKPERRLTIEGVLDHPWLNSTEALDNVLPQAQLMMDKAVVAGIQQAHA  
40 EQLANMRIQDLKVSLLKPLHSVNNPILRKRKLLGTPKDSVYIHDHENGAEEDSNVALEK  
LRDVIAQCILPQAGENEDEKLNEVMQEAWKYNRECKLLRDTLQSFSWNGRGFTDKVDR  
LKLAEIVKQVIEEQTTSHESQ"

BASE COUNT 571 a 445 c 485 g 467 t

45 ORIGIN

1 ccagcctaga gccgcccggc gaagcagagc cggcgccggg gtcctcatcc ccaccggtcc  
61 cgaggggccc ctgctgcccg tcgccacgag gcccaggggc ccgagtgccg agccctttgc  
121 tccctcggcc gcgcggggac agggctgctg agcagcctcc gcctctcccg gctgtggggg  
181 cccactgag tatgtcggag gagagcgaca tggacaaagc catcaaggaa acttccattt  
50 241 tagaagaata cagtatcaat tggactcaga agctgggagc tggaattagt ggtccagtta  
301 gagtctgtgt aaagaaatct actcaagaac ggtttgcgct gaaaattctt cttgatcgtc  
361 caaaagctag aaatgaggta cgtctgcaca tgatgtgtgc cacacacca aacatagttc  
421 agattattga agtgtttgct aacagtgtcc agtttcccca tgagtccagc cctagggccc  
481 gactcttaat tgtaatggag atgatggaag ggggagagct atttcacaga atcagccagc  
55 541 accggcactt tacagagaag caagccagcc aagtaacaaa gcagatagct ttggctctgc  
601 ggcactgtca cttgttaaac attgcgcaca gagacctcaa gcctgaaaat ctgcttttta  
661 aggataactc tttggatgcc ccagtgaagt tgtgtgactt tggatttgcc aagattgacc  
721 aaggtgactt gatgacaccc cagttcaccc cttattatgt agcaccacag gtactggagg  
781 cgcaaagaag gcatcagaag gagaaatctg gcatcatacc tacctcaccg acgccctaca  
60 841 cttacaacaa gagctgtgac ttgtgggtccc taggggtgat tatctatgtg atgctgtgcg  
901 gataccctcc tttttactcc aaacaccaca gccggactat cccaaaggat atgcgaagaa  
961 agatcatgac aggcagtttt gacttcccag aggaagagtg gagtcatgac tcagagatgg  
1021 ccaaagatgt tgtgaggaag ctctgaagg tcaaacggga gaggagactc accatcgagg  
1081 gagtgtctgga ccaccctgg ctcaattcca ccgaggccct ggataatgtg ctgccttctg



1141 ctcagctgat gatggacaag gcagtggttg caggaatcca gcagggtcac gcggaacagt  
1201 tggccaacat gagaatccag gatctgaaag tcagcctcaa acccctgcac tcagtgaaca  
1261 accccattct gcggaagagg aagttacttg gcaccaagcc aaaggacagt gtctatatcc  
1321 acgaccatga gaatggagcc gaggattcca atgttgccct ggaaaaactc cgagatgtga  
5 1381 ttgctcagtg tattctcccc caggctggag agaatgaaga tgagaaactg aatgaagtaa  
1441 tgcaggaggc ttggaagtat aaccgggaat gcaaactcct aagagatact ctgcagagct  
1501 tcagctggaa tggtcgtgga ttcacagata aagtagatcg actaaaactg gcagaaattg  
1561 tgaagcaggt gatagaagag caaaccacgt cccacgaatc ccaataatga cagcttcaga  
1621 ctttgttttt ttaacaattt gaaaaattat tctttaatgt ataaagtaat tttatgtaaa  
10 1681 ttaataaatc ataatttcac ttccacattg attaaagctg ctgtatagat ttagggtgca  
1741 ggacttaata atagtatagt tattgtttgt ttttaagaaa agctcagttc tagagacata  
1801 ctattacttt aggactgtgt agttgtatat ttgtaagatg acagatgatg ctgtcaagca  
1861 atattgtttt atttgtaata aaatatacaa aaatcacttg ccagcagtag aaaaaggacc  
1921 gactataccg acctttctga ttagtaaaca gttgaatcaa ggactctg  
15

20 DEFINITION Human insulin-stimulated protein kinase 1 (ISPK-1) mRNA,  
complete

cds.

ACCESSION U08316

VERSION U08316.1 GI:475587

FEATURES Location/Qualifiers

25 source

1..2260

/organism="Homo sapiens"

/db\_xref="taxon:9606"

/chromosome="X"

/map="Xp21.3-22.1"

30 /clone\_lib="MOLT 4 T-cell leukemia and human

placenta cDNA

libraries"

gene

1..2223

/gene="ISPK-1"

35 CDS

1..2223

/gene="ISPK-1"

/note="ribosomal protein S6 kinase"

/codon\_start=1

/product="insulin-stimulated protein kinase 1"

40 /protein\_id="AAA81952.1"

/db\_xref="GI:475588"

/translation="MPLAQLADPWQKMAVESPSDSAENGQQIMDEPMGEEEEINPQTEE

45 VSIKEIAITHHVKEGHEKADPSQFELLKVLGQGSFGKVFLVKKISGSDARQLYAMKVL

KKATLTKVRDRVRTKMERDILVEVNHPFIVKLHYAFQTEGKLYLILDFLRGGDLFTRL

50 KEVMFTEEDVKFYLAELALALDHLHSLGIIYRDLKPENILLDEEGHIKLTDFGLSKES

IDHEKKAYSFCGTVEYMAPEVVNRRGHTQSADWWSFGVLMFEMLTGTLPFQGKDRKET

MTMILKAKLGMPQFLSPEAQSLRLMLFKRNPANRLGAGPDGVEEIKRHSFFSTIDWNK

55 LYRREIHPPFKPATGRPEDTFYFDPEFTAKTPKDSPIPPSANAHQLFRGFSFVAITS

DDESQAMQTVGVHSIVQQLHRNSIQFTDGYEVKEDIGVGSYSVCKRCIHKATNMEFAV

60 KIIDKSKRDPTEIEILLRYGQHPNIITLKD VYDDGKYVYVVT ELMKGGELLDKILRQ

KFFSEREASAVLFTITKTVEYLHAQGVVHRDLKPSNILYVDESGNPESIRICDFGFAK

QLRAENGLLMTPCYTANFVAPEVLKRQGYDAACDIWSLGVLLYTMLTGYTPFANGPDD

TPEEILARIGSGKFSLSGGYWNSVSDTAKDLVSKMLHVDPHQRLTAALVLRHPWIVHW

DQLPQYQLNRQDAPHLVKGAMAATYSALNRNQSPVLEPVGRRSTLAQRRGIKKITSTAL

5

BASE COUNT            704 a       417 c       510 g       629 t  
ORIGIN

```

1  atgccgctgg  cgcagctggc  ggaccctggg  cagaagatgg  ctgtggagag  cccgtccgac
61  agcgctgaga  atggacagca  aattatggat  gaacctatgg  gagaggagga  gattaaccca
10 121  caaactgaag  aagtcagtat  caaagaaatt  gcaatcacac  atcatgtaaa  ggaaggacat
181  gaaaaggcag  atccttccca  gtttgaactt  ttaaagatat  tagggcaggg  atcatttgga
241  aagggttttct  tagttaaaaa  aatctcaggg  tctgatgcta  ggcagcttta  tgccatgaag
301  gtattgaaga  aggccacact  gaaagttcga  gaccgagttc  ggacaaaaat  ggaacgtgat
361  atcttggtag  aggttaatca  tccttttatt  gtcaagttgc  attatgcttt  tcaaactgaa
15 421  gggaagttgt  atcttatttt  ggattttctc  aggggaggag  atttgtttac  acgcttatcc
481  aaagaggtga  tgttcacaga  agaagatgtc  aaattctact  tggctgaact  tgcacttgct
541  ttagaccatc  tacatagcct  gggaataatt  tatagagact  taaaaccaga  aaatatactt
601  cttgatgaag  aaggtcacat  caagttaaca  gatttcggcc  taagtaaaga  gtctattgac
661  catgaaaaga  aggcattatc  tttttgtgga  actgtggagt  atatggctcc  agaagtagtt
20 721  aatcgctcag  gtcatactca  gagtgtgcac  tgggtgtcct  ttggtgtggt  aatgtttgaa
781  atgcttactg  gtacactccc  tttccaagga  aaagatcgaa  aagaaacaat  gactatgatt
841  cttaaagcca  aacttggaa  gccacagttt  ttgagtcctg  aagcgcagag  tcttttacga
901  atgcttttca  agcgaatatc  tgcaaacaga  ttaggtgcag  gaccagatgg  agttgaagaa
961  attaaaagac  attcattttt  ctcaacgata  gactggaata  aactgtatag  aagagaaatt
25 1021  catccgccat  ttaaacctgc  aacgggcagg  cctgaagata  cattctattt  tgatcctgag
1081  tttactgcaa  aaactcccaa  agattcacct  ggcattccac  ctagtgtcaa  tgcacatcag
1141  ctttttcggg  ggtttagtgt  tgttgctatt  acctcagatg  atgaaagcca  agctatgcag
1201  acagttagtg  tacattcaat  tgttcagcag  ttacacagga  acagtattca  gtttactgat
1261  ggatatgaag  taaaagaaga  tattggagtt  ggctcctact  ctgtttgcaa  gagatgtata
30 1321  cataaagcta  caaacatgga  gtttgacagt  aagattattg  ataaaagcaa  gagagacca
1381  acagaagaaa  ttgaaattct  tcttcgttat  ggacagcatc  caaacattat  cactctaaag
1441  gatgtatatg  atgatggaaa  gtatgtgtat  ttagtaacag  aacttatgaa  aggagtgaa
1501  ttgctggata  aaattcttag  acaaaaattt  ttctctgaac  gagaggccag  tgctgtcctg
1561  tttactataa  ctaaaaccgt  tgaatatctt  cagcacaag  ggggtggttca  tcgagacttg
35 1621  aaacctagca  acattcttta  tgtggatgaa  tctggtaatc  cggaatctat  tcgaatttgt
1681  gattttggct  ttgcaaaaca  gctgagagcg  gaaaatggtc  ttctcatgac  tccttggtac
1741  actgcaaatt  ttgttgcaac  agaggtttta  aaaagacaag  gctatgatgc  tgcttgatg
1801  atatggagtc  ttggtgtcct  actctataca  atgcttaccg  gttacactcc  atttgcaaat
1861  ggtcctgatg  atacaccaga  ggaaatattg  gcacgaatag  gtagcggaaa  attctcactc
40 1921  agtggtggtt  actggaattc  tgtttcagac  acagcaaagg  acctggtgtc  aaagatgctt
1981  catgtagacc  ctcatcagag  actgactgct  gctcttggtc  tcagacatcc  ttggatcgtc
2041  cactgggacc  aactgccaca  ataccaacta  aacagacagg  atgcaccaca  tctagtaaag
2101  ggtgccatgg  cagctacata  ttctgctttg  aaccgtaatc  agtcaccagt  tttggaacca
2161  gtaggccgct  ctactcttgc  tcagcggaga  ggtattaaaa  aaatcacctc  aacagccctg
45 2221  tgaagtgacc  tcagttagat  atttggatcc  atgggtgtaa

```

50 DEFINITION Homo sapiens discoidin domain receptor family, member 1 (DDR1),

transcript variant 3, mRNA.

ACCESSION NM\_013994

VERSION NM\_013994.1 GI:7669484

55 EC number = "2.7.1.112"

60

DEFINITION H.sapiens mRNA for protein kinase C mu.

ACCESSION X75756

VERSION X75756.1 GI:438372

FEATURES Location/Qualifiers

source 1..3742  
/organism="Homo sapiens"  
/db\_xref="taxon:9606"  
/tissue\_type="placenta"  
5 CDS 236..2974  
/codon\_start=1  
/product="protein kinase C mu"  
/protein\_id="CAA53348.1"  
/db\_xref="GI:438373"  
10 /db\_xref="SWISS-PROT:Q15139"  
/translation="MSAPPVLRPPSPLLPVAAAAAAAAAALVPGSGPGPAPFLAPVAA  
PVGGISFHLQIGLSREPVLLLQDSSGDYSLAHVREMACSIVDQKFPECGFYGMYDKIL  
15 LFRHDPTSENILQLVKAASDIQEGDLIEVVLSRSATFEDFQIRPHALFVHSYRAPAFC  
DHCGEMLWGLVRQGLKCEGCGLNYHKRCAFKIPNNCSGVRRRRLSNVSLTGVSTIRTS  
20 SAELSTSAPDEPLLQKSPSEFIGREKRSNSQSYIGRPIHLDKILMSKVVPHTFVIH  
SYTRPTVCQYCKLLKGLFRQGLQCKDCRFNCHKRCAPKVPNNCLGEVTINGDLLSPG  
AESDVVMEEGSDDNDSENRSGLMDDMEEAMVQDAEMAMAECQNDSGEMQDPDPDHEDA  
25 NRTISPSTSNNIPLMRVVQSVKHTKRKSSTVMKEGWMVHYTSKDTLRKRHYWRLDSKC  
ITLQNDTGSRYKEIPLSEILSLEPVKTSALIPNGANPHCFEITTANVVVYVGENVV  
30 NPSSPSPNNSVLTSVGADVARMWEIAIQHALMPVIPKGSSVGTGTNLHRDISVSISV  
SNCQIQENVDISTVYQIFPDEVLGSGQFGIVYGGKHRKTGRDVAIKIIDKLRFPTKQE  
SQLRNEVAILQNLHHPGVVNLECMFETPERVFVMEKLGDMLEMILSSEKGRLEPHI  
35 TKFLITQILVALRHLHFKNIVHCDLKPENVLLASADPFQVKLCDFGFARIIGESFR  
RSVVGTPAYLAPEVLRNKGYNRSLDMWSVGVIIYVSLSGTFPFNEDEDIHDQIQNAAF  
40 MYPPNPWKEISHEAIDLINLLQVKMRKRYSDKTLSPWLQDYQTWLRLRELECKIG  
ERYITHESDDLWEKYAGEQRLQYPHTLINPSASHSDTPETEEMKALGERVSIL"  
BASE COUNT 1015 a 897 c 865 g 965 t

## 45 ORIGIN

1 gaattccttc tctcctcctc ctgcgccttc tctcgccttc cctcctcctc ctgcgccttc  
61 cctcccgatc ctcatccctc tgccctcccc cagcccaggg acttttcggg aaagttttta  
121 ttttccgtct gggctctcgg agaaagaagc tcttggtca gcggctgcaa aactttcctg  
181 ctgccgcgcc gccagcccc gccctccgct gcccgccct gcgccccgcc gagcgatgag  
50 241 cgccctccg gtcctgcggc cgccagtcg gctgctgccc gtggcgccgg cagctgccgc  
301 agcgcccgcc gactgggtcc cagggtccgg gcccgggccc gcgcccgttc tggctcctgt  
361 cgcggccccc gtcgggggca tctcgttcca tctgcagatc ggcctgagcc gtgagccggt  
421 gctgctgtcg caggactcgt ccgggggacta cagcctggcg cagctccgag agatggcttg  
481 ctccattgtc gaccagaagt tccctgaatg tggtttctac ggaatgtatg ataagatcct  
55 541 gctttttcgc catgacccta cctctgaaaa catccttcag ctggtgaaag cggccagtga  
601 tatccaggaa ggcgatctta ttgaagtggc cttgtcacgt tccgccacct ttgaagactt  
661 tcagattcgt cccacgctc tctttgttca ttcatacaga gtcacagctt tctgtgatca  
721 ctgtggagaa atgctgtggg ggctggtacg tcaaggtctt aaatgtgaag ggtgtggtct  
781 gaattaccat aagagatgtg catttaaaat acccaacaat tgcagcggtg tgaggcggag  
60 841 aaggctctca aacgtttccc tcaactgggg cagcaccatc cgcacatcat ctgctgaact  
901 ctctacaagt gccctgatg agcccttctt gcaaaaatca ccatcagagt cgtttattgg  
961 tcgagagaag aggtcaaatt ctcaatcata cattggacga ccaattcacc ttgacaagat  
1021 tttgatgtct aaagttaaag tgccgcacac atttgtcatc cactcctaca cccggccac  
1081 agtgtgccag tactgcaaga agcttctgaa ggggcttttc aggcagggct tgcagtgcac

```

1141 agattgcaga ttcaactgcc ataaacgttg tgcaccgaaa gtaccaaaaca actgccttgg
1201 cgaagtgacc attaatggag atttgcttag ccctggggca gagtctgatg tggatcatgga
1261 agaaggagat gatgacaatg atagtgaag gaacagtggg ctcattggatg atattggaaga
1321 agcaatgggtc caagatgcag agatggcaat ggcagagtgc cagaacgaca gtggcgagat
5 1381 gcaagatcca gaccagacc acgaggacgc caacagaacc atcagtccat caacaagcaa
1441 caatatccca ctcattgaggg tagtgcagtc tgtcaaacac acgaagagga aaagcagcac
1501 agtcatgaaa gaaggatgga tgggtccacta caccagcaag gacacgctgc ggaaacggca
1561 ctattggaga ttggatagca aatgtattac cctctttcag aatgacacag gaagcaggtg
1621 ctacaaggaa attcctttat ctgaaatttt gtctctggaa ccagtaaaaa cttcagcttt
10 1681 aattcctaatt ggggccaatc ctcattgttt cgaaatcact acggcaaatg tagtgtatta
1741 tgtggggagaa aatgtggtca atccttccag cccatcacca aataacagtg ttctcaccag
1801 tggcggttggg gcagatgtgg ccaggatgtg ggagatagcc atccagcatg cccttatgcc
1861 cgtcattccc aagggtcctt ccgtgggtac aggaaccaac ttgcacagag atatctctgt
1921 gagtattttca gtatcaaatt gccagattca agaaaatgtg gacatcagca cagtatatca
15 1981 gatttttcct gatgaagtac tgggttctgg acagtttggg attgtttatg gaggaaaaa
2041 tcgtaaaaaa ggaagagatg tagctattaa aatcattgac aaattacgat ttccaacaaa
2101 acaagaaagc cagcttcgta atgaggttgc aattctacag aaccttcac accctgggtg
2161 tgtaaatattg gagtgtatgt ttgagacgcc tgaaagagtg tttgttgtta tggaaaaact
2221 ccatggagac atgctggaaa tgatcttgtc aagtgaanaa ggcaggttgc cagagcacat
20 2281 aacgaagttt ttaattactc agatactcgt ggctttgcgg caccttcatt ttaaaaatat
2341 cgttccactgt gacctcaaac cagaaaatgt gttgctagcc tcagctgac cttttcctca
2401 ggtgaaactt tgtgattttg gttttgcccg gatcattgga gagaagtctt tccggaggtc
2461 agtgggtgggt acccccgctt acctggctcc tgaggtccta aggaacaagg gctacaatcg
2521 ctctctagac atgtggtctg ttggggtcat catctatgta agcctaagcg gcacattccc
25 2581 atttaatgaa gatgaagaca tacacgacca aattcagaat gcagctttca tgtatccacc
2641 aaatccctgg aaggaaatat ctcattgaagc cattgatctt atcaacaatt tgcgtcaagt
2701 aaaaatgaga aagcgctaca gtgtggataa gaccttgagc cacccttggc tacaggacta
2761 tcagacctgg ttagatttgc gagagctgga atgcaaaatc ggggagcgtc acatcaccca
2821 tgaaagtgat gacctgaggt gggagaagta tgcaggcgag cagcggctgc agtaccacac
30 2881 acacctgatc aatccaagtg ctagccacag tgacactcct gagactgaag aaacagaaat
2941 gaaagccctc ggtgagcgtg tcagcatcct ctgagttcca tctcctataa tctgtcaaaa
3001 cactgtggaa ctaataaata catacggtca ggtttaacat ttgccttgca gaactgccat
3061 tattttctgt cagatgagaa caaagctggt aaactgttag cactgttgat gtatctgagt
3121 tgccaagaca aatcaacaga agcattttgta ttttgtgtga ccaactgtgt tgtattaaca
35 3181 aaagttccct gaaacacgaa acttgttatt gtgaatgatt catgttatat ttaatgcatt
3241 aaacctgtct ccactgtgcc tttgcaaatc agtgttttct ttactggagc ttcatttttg
3301 taagagacag aatgtatctg tgaagtagtt ctgtttgggtg tgtcccattg gtgtgtcat
3361 tgtaaacaaa ctcttgaaga gtcgattatt tccagtgttc tatgaacaac tccaaaaccc
3421 atgtgggaaa aaaatgaatg aggaggttag ggaataaaat cctaagacac aaatgcatga
40 3481 acaagtttta atgtatagtt ttgaaatcctt tgcctgcctg gtgtgcctca gtatatttaa
3541 actcaagaca atgcacctag ctgtgcaaga cctagtgtct ttaagcctaa atgccttaga
3601 aatgtaaaact gccatatata acagatacat ttccctcttt cttataatac tctgttgtac
3661 tatggaaaat cagctgctca gcaacctttc acctttgtgt atttttcaat aataaaaaat
3721 attcttgtca aaaaaaaaaa aa
45

```

```

50 DEFINITION Human protein kinase C-theta (PRKCT) mRNA, complete cds.
ACCESSION L01087
VERSION L01087.1 GI:558098
FEATURES
    source 1..2754
           /organism="Homo sapiens"
55         /db_xref="taxon:9606"
           /cell_line="HEL"
           /cell_type="erythroleukemia"
           /tissue_lib="HEL cell lambda-gt11 library of M.
60 Poncz"
    gene 95..2215
           /gene="PRKCT"
    CDS 95..2215
           /gene="PRKCT"
           /codon_start=1

```

/product="protein kinase C-theta"  
 /protein\_id="AAA75571.1"  
 /db\_xref="GI:558099"

5 /translation="MSPFLRIGLSNFDGSCQSCQGEAVNPYCAVLVKEYVESENGQM  
 YIQKKPTMYPPWDSTFDAHINKGRVMQIIVKGKNVDLISETTVELYSLAERCCKNNKG  
 TEIWLELKPQGRMLMNARYFLEMSDTKDMNEFETEGFFALHQRRGAIKQAKVHHVKCH  
 10 EFTATFFPQPTFCSVCHEFVWGLNKQGYQCRQCNAAIHKKCIDKVIKCTGSAINSRE  
 TMFHKERFKIDMPHRFKVYNYKSPTFCEHCGLLWGLARQGLKCDACGMNVHHRQCQTK  
 15 VANLCGINQKLMAEALAMIESTQQARCLRDTEQIFREGPVEIGLPCSIKNEARLPCLP  
 TPGKREPQGISWESPLDEVDKMCHLPEPELNKERPSLQIKLKIEDFILHKMLGKGSFG  
 KVFLAEFKKTNQFFAIKALKKDVLMDDDVECTMVEKRVLSLAWEHPFLTHMFCTFQT  
 20 KENLFFVMEYLNCGDLMYHIQSCHKFDLSRATFYAAEIIILGLQFLHSGIVYRDLKLD  
 NILLDKDGHIKIADFGMCKENMLGDAKTNTFCGTPDYLAPEILLGQKYNHSDVWSFG  
 25 VLLYEMLIGQSPFHGQDEEELFHSIRMDNPFYPRWLEKEAKDLLVKLFVREPEKRLGV  
 RGDIRQHPLFREINWHEELERKEIDPPFRPKVKS PFDCSNFDKEFLNEKPRLSFADRAL  
 INSMQNMFRNFSFMNPRMERLIS"  
 BASE COUNT 764 a 657 c 686 g 647 t  
 30 ORIGIN  
 1 gaattccgcc agccccgccca gtccccgcgc agtccccgcg cagtcccagc gccaccgggc  
 61 agcagcggcg cegtgtctgc tccagggcgc aaccatgtcg ccatttcttc ggattggctt  
 121 gtccaacttt gactgcggt cctgccagtc ttgtcagggc gaggctgtta acccttactg  
 181 tgctgtgctc gtcaaagagt atgtcgaatc agagaacggg cagatgtata tccagaaaaa  
 35 241 gcctaccatg taccaccct gggacagcac ttttgatgcc catatcaaca agggagaggt  
 301 catgcagatc atttgtgaaag gcaaaaacgt ggacctcatc tctgaaacca ccgtggagct  
 361 ctactcgctg gctgagaggt gcaggaagaa caacgggaag acagaaatat ggttagagct  
 421 gaaacctcaa ggccgaatgc taatgaatgc aagatacttt ctggaaatga gtgacacaaa  
 481 ggacatgaat gaatttgaga cggaaggctt ctttgccttg catcagcgcc ggggtgccat  
 40 541 caagcaggca aaggtccacc acgtcaagtg ccacgagttc actgccacct tctcccaca  
 601 gcccacattt tgctctgtct gccacgagtt tgtctggggc ctgaacaaac agggctacca  
 661 gtgccgacaa tgcaatgcag caattcacaa gaagtgtatt gataaagtta tagcaaagtg  
 721 cacaggatca gctatcaata gccgagaaac catgttccac aaggagagat tcaaaattga  
 781 catgccacac agatttaaag tctacaatta caagagcccg accttctgtg aacactgtgg  
 45 841 gaccctgtcg tggggactgg cacggcaagg actcaagtgt gatgcattgt gcatgaatgt  
 901 gcatcataga tgccagacaa aggtggccaa cctttgtggc ataaaccaga agctaattgc  
 961 tgaagcgctg gccatgattg agagcactca acaggctcgc tgcttaagag atactgaaca  
 1021 gatcttcaga gaaggtccgg ttgaaattgg tctcccatgc tccatcaaaa atgaagcaag  
 1081 gctgccatgt ttaccgacac cgggaaaaag agagcctcag ggcatttcct gggagtctcc  
 50 1141 gttggatgag gtggataaaa tgtgccatct tccagaacct gaactgaaca aagaaagacc  
 1201 atctctgcag attaaactaa aaattgagga ttttatcttg cacaaaatgt tggggaaagg  
 1261 aagttttggc aaggtcttcc tggcagaatt caagaaaacc aatcaatttt tcgcaataaa  
 1321 ggccttaaag aaagatgtgg tcttgatgga cgatgatgtt gagtgcacga tggtagagaa  
 1381 gagagttctt tccttggcct gggagcatcc gtttctgacg cacatgtttt gtacatttca  
 55 1441 gaccaaggaa aacctctttt ttgtgatgga gtacctcaac ggaggggact taatgtacca  
 1501 catccaaagc tgccacaagt tcgaccttcc cagagcgacg ttttatgctg ctgaaatcat  
 1561 tcttggctcg cagttccttc attccaaagg aatagtctac agggacctga agctagataa  
 1621 catcctgcta gacaaagatg gacatatcaa gatcgcgat tttggaatgt gcaaggagaa  
 1681 catgttagga gatgccaaaga cgaataacct ctgtgggaca cctgactaca tcgccccaga  
 60 1741 gatcttgctg ggtcagaaat acaaccactc tgtggactgg tggctcctcg gggttctcct  
 1801 ttatgaaatg ctgattggtc agtcgccttt ccacgggcag gatgaggagg agctcttcca  
 1861 ctccatccgc atggacaatc ccttttacct acggtggctg gagaagggaag caaaggacct  
 1921 tctgggtgaag ctcttcgtgc gagaacctga gaagaggctg ggcgtgaggg gagacatccg  
 1981 ccagcaccct ttgtttcggg agatcaactg ggaggaactt gaacggaagg agattgacct

```
2041 accgttccgg ccgaaagtga aatcaccatt tgactgcagc aatttcgaca aagaattctt
2101 aaacgagaag ccccggtgt catttgccga cagagcactg atcaacagca tggaccagaa
2161 tatgttcagg aacttttcct tcatgaaccc ccgatggag cggctgatat cctgaatctt
2221 gcccctccag agacaggaaa gaatttgcct tgtccctggg aactggttca agagacactg
5 2281 cttgggttcc tttttcaact tggaaaaaga aagaaacact caacaataaa gactgagacc
2341 cggttcgccc catgtgactt ttatctgtag cagaaaccaa gtctacttca ctaatgacga
2401 tgccgtgtgt ctgctctcct gacatgtctc acagacgctc ctgaagttag gtcattacta
2461 accatagtta tttacttgaa agatgggtct ccgacttgg aaaggtttca agacttgata
2521 ctgcaataaa ttatggctct tcacctgggc gccaaactgct gatcaacgaa atgcttggtg
10 2581 aatcaggggc aaacggagta cagacgtctc aagactgaaa cggccccatt gcctggtcta
2641 gtagcggtc tcaactcagc gcagacaagt aatcactaac ccgttttatt ctattcctat
2701 ctgtggatgg gtaaatgctg ggggccagcc ctggataggt ttttatggga attc
```

15

DEFINITION Homo sapiens mRNA for AMP-activated protein kinase beta 2 subunit.

ACCESSION AJ224538

20 VERSION AJ224538.1 GI:2916801

FEATURES Location/Qualifiers

source

1..819

/organism="Homo sapiens"

/db\_xref="taxon:9606"

25 gene

1..819

/gene="AMPK beta 2"

CDS

1..819

/gene="AMPK beta 2"

/function="regulatory subunit"

30 /codon\_start=1

/evidence=experimental

/product="AMP-activated protein kinase beta 2

subunit"

/protein\_id="CAA12030.1"

35 /db\_xref="GI:2916802"

/db\_xref="SWISS-PROT:O43741"

/translation="MGNTTSDRVSGERHGAKAARSEGAGGHAPGKEHKIMVGSTDDPS

40 VFSLPDSKLPDKEFVSWQQDLEDSVKPTQQARPTVIRWSEGGKEVFISGSFNNWSTK

IPLIKSHNDFVAILDLPEGEHQYKFFVDGQVHDPSEPVVTSQLGTINNLIHVKKSDF

EVFDALKLDSMESSETSCRDLSSSPGPYGQEMYAFRSEERFKSPPIPPHLLQVILN

45 KDTNISCDPALLPEPNHVMLNHLIALSIKDSVMVLSATHRYKKKYVTTLLYKPI"

BASE COUNT 208 a 215 c 197 g 199 t

ORIGIN

```
50 1 atgggaaaca ccaccagcga cggggtgtcc ggggagcgcc acggcgccaa ggctgcacgc
61 tccgagggcg caggcgcca tgcccgggg aaggagcaca agatcatggt ggggagtacg
121 gacgaccca gcggtgtcag cctccctgac tccaagctcc ctggggacaa agagtgtgta
181 tcatggcagc aggatttggg ggactccgta aagccacac agcaggcccg gccactgtt
241 atccgctggt ctgaaggagg caaggaggtc ttcatctctg ggtctttcaa caattggagc
55 301 accaagattc cactgattaa gagccataat gactttgttg ccatcctgga cctccctgag
361 ggagagcacc aatacaagtt ctttgtggat ggacagtggg ttcatgatcc atcagagcct
421 gtggttacca gtcagcttgg cacaattaac aatttgatcc atgtcaagaa atctgatttt
481 gaggtgttcg atgcttttaa gttagattct atggaaagt ctgagacatc ttgtagagac
541 ctttccagct cacccccagg gccttatggt caagaaatgt atgcgtttcg atctgaggaa
60 601 agattcaaat cccaccccat ccttctcct catctacttc aagttattct taacaaagac
661 actaatatatt cttgtgaccc agccttactc cctgagccca accatgttat gctgaacat
721 ctctatgcat tgtccattaa ggacagtgtg atggtcctta gcgcaaccca tcgctacaag
781 aagaagtatg ttactactct gctatacaag cccatttga
```

DEFINITION Human protein kinase (JNK2) mRNA, complete cds.

5 ACCESSION U09759  
VERSION U09759.1 GI:607785

FEATURES  
source Location/Qualifiers  
1..1873  
/organism="Homo sapiens"  
10 /db\_xref="taxon:9606"  
gene 152..1426  
/gene="JNK2"  
CDS 152..1426  
/gene="JNK2"  
15 /codon\_start=1  
/product="protein kinase"  
/protein\_id="AAA74740.1"  
/db\_xref="GI:607786"

20 /translation="MSDSKCDSQFYSVQVADSTFTVLKRYQQLKPIGSGAQGIVCAAF  
DTVLGINVAVKKLSRPFQNQTHAKRAYRELVLLKCVNHKNIISLLNVFTPOKTLLEEFQ  
DVYLVMEMLMDANLCQVIHMELDHERMSYLLYQMLCGIKHLHSAGIIHRDLKPSNIVVK  
25 SDCTLKILDFGLARTACTNFMTPYVVTRYRAPEVILGMGYKENVDIWSVGCIMGEL  
VKGCVIFQGTDHIDQWNKVIEQLGTPSAEFMKKLQPTVRNYVENRPKYPGIKFEELFP  
30 DWIFPSESERDKIKTSQARDLLSKMLVIDPDKRISVDEALRHPYITVWYDPAEAEAPP  
PQIYDAQLEEREHAIEEWKELIYKEVMDWEERSKNGVVKDQPPDAAVSSNATPSQSSS  
INDISSMSTEQTLASDSTDSSLDASTGPLEGCR"

BASE COUNT 562 a 388 c 427 g 496 t

35 ORIGIN  
1 caaactacgt gctgtacagc tgcatacagc gctcgtagac atgtccagca gctgggtcgag  
61 gtccacgccg cggtaggtga agttgcccga ggtccggcga gggatctgaa acttgcccct  
121 tacccttcgg gatattgcag gacgctgcat catgagcgac agtaaatgtg acagtcagtt  
181 ttatagtgtc caagtggcag actcaacctt cactgtccta aaacggtacc agcagctgaa  
40 241 accaattggc tctggggccc aagggattgt ttgtgctgca tttgatacag ttcttgggat  
301 aaatggtgca gtcaagaaac taagccgtcc ttttcagaac caaactcatg caaagagagc  
361 ttatcgtgaa cttgtcctct taaaatgtgt caatcataaa aatataatta gtttgttaaa  
421 tgtgtttaca ccacaaaaaa ctctagaaga atttcaagat gtgtatttgg ttatggaatt  
481 aatggatgct aacttatgtc aggttattca catggagctg gatcatgaaa gaatgtccta  
45 541 ccttcctttac cagatgcttt gtggtattaa acatctgcat tcagctggta taattcatag  
601 agatttgaag cctagcaaca ttggttgtaa atcagactgc accctgaaga tccttgactt  
661 tggcctggcc cggacagcgt gcactaactt catgatgacc ccttacgtgg tgacacggta  
721 ctaccgggcg cccgaagtca tcctgggtat gggctacaaa gagaacgttg atatctggtc  
781 agtgggttgc atcatgggag agctggtgaa aggttgtgtg atattccaag gcactgacca  
50 841 tattgatcag tggaataaag ttattgagca gctgggaaca ccatcagcag agttcatgaa  
901 gaaacttcag ccaactgtga ggaattatgt cgaaaacaga ccaaagtatc ctggaatcaa  
961 atttgaagaa ctctttccag attggatatt cccatcagaa tctgagcgag acaaaataaa  
1021 aacaagtcaa gccagagatc tgttatcaaa aatgttagtg attgatcctg acaagcggat  
1081 ctctgtagac gaagctctgc gtcacccata catcactgtt tggatagacc ccgccgaagc  
55 1141 agaagcccca ccacctcaa tttatgatgc ccagttggaa gaaagagaac atgcaattga  
1201 agaatggaaa gagctaattt acaaagaagt catggattgg gaagaaagaa gcaagaatgg  
1261 tgttgtaaaa gatcagcctc cagatgcagc agtaagtagc aacgccactc cttctcagtc  
1321 ttcatcgatc aatgacattt catccatgct cactgagcag acgctggcct cagacacaga  
1381 cagcagtcct gatgcctcga cgggacccct tgaaggctgt cgatgatagg ttagaaatag  
60 1441 caaacctgtc agcattgaag gaactctcac ctccgtgggc ctgaaatgct tgggagttga  
1501 tggaaccaa tagaaaaact ccatgttctg catgtaagaa acacaatgcc ttgccctact  
1561 cagacctgat aggattgcct cgttagatga taaaatgagg cagaatatgt ctgaagaaaa  
1621 aaattgcaag ccacacttct agagattttg ttcaagatca tttcagttga gcagttagag  
1681 taggtgaatt tgtcaaattg tactagtgc agtttctcat catctgtaac tgttgagatg

9/17

1741 attgtgcatg tgaccacaaa tgcttgcttg gacttgccca tctagcactt tggaaatcag  
1801 tattttaaatg ccaaataatc ttccaggtag tgctgcttct gaagtatatct cttaatcctc  
1861 ttaagtaatt tgg

5

DEFINITION Human p21-activated protein kinase (Pak2) gene,  
complete cds.

10 ACCESSION U24153  
VERSION U24153.1 GI:780807  
FEATURES Location/Qualifiers  
source 1..2019  
/organism="Homo sapiens"  
15 /db\_xref="taxon:9606"  
/clone="212"  
/cell\_type="EBV-transformed peripheral lymphocytes"  
/clone\_lib="lambdaYESR human lymphocyte cDNA  
library;  
20 Elledge and Spottswood, EMBO 10:2653(1991)"  
gene 40..1617  
/gene="Pak2"  
CDS 40..1617  
/gene="Pak2"  
25 /codon\_start=1  
/product="p21-activated protein kinase"  
/protein\_id="AAA65442.1"  
/db\_xref="GI:780808"  
30 /translation="MSDNGELEDKPPAPPVRMSSTIFSTGGKDPLSANHSLKPLPSVP  
EEKKPRHKIISIFSGTEKSGSKKKEKERPEISPPSDFEHTIHVGFD AVTGEFTGMPEQW  
ARLLQTSNITKLEQKKNPQAVLDVLKFYDSNTVKQKYLSTTPPEKDGLPSGTPALNAK  
35 GTEAPAVVTEEEDDDEETAPPVIAPRPDHTKSIYTRSVIDPVPAPVGDSHVDGAAKSL  
DKQKKKPKMTDEEIMEKLRTIVSIGDPKKKYTRYEKIGQGASGTVFTATDVALGQEV  
40 IKQINLQKQPKKELINEILVMKELKNPNIVNFLDSYLVGDEL FVVM EYLAGGSLTDV  
VTETACMDEAQIAAVCRECLQALEFLHANQVIHRDIKSDNVLLGMEGSVKLTDGFCA  
QITPEQSKRSTMVGTPTYWMAPEVVTRKAYGPKVDIWSLGIMAIEMVEGEPPYLNENPL  
45 RALYLIATNGTPELQNPKEKLSPIFRDFLNRCLMDVEKRGSAKELLQHPFLKLAKPLS  
SLTPLIMAAKEAMKSNR"  
BASE COUNT 615 a 413 c 481 g 510 t

50 ORIGIN

1 gaccttggct tgcccggggc catttcataa ttctgaatca tgtctgataa cggagaactg  
61 gaagataagc ctccagcacc tcctgtgcga atgagcagca ccatcttttag cactgggaggc  
121 aaagaccctt tgtcagccaa tcacagtttg aaacctttgc cctctgttcc agaagagaaa  
181 aagcccaggc ataaaatcat ctccatattc tcaggcacag agaaaggaag taaaaagaaa  
55 241 gaaaaggaac ggccagaaat ttctcctcca tctgattttg agcacaccat ccatgttggc  
301 tttgatgctg ttactggaga attcactggc atgccagaac agtggggtcg attactacag  
361 acctccaata tcaccaaact agagcaaaag aagaatcctc aggtgtgtgc ggatgtccta  
421 aagttctacg actccaacac agtgaagcag aaatatctga gctttactcc tcctgagaaa  
481 gatggccttc cttctggaac gccagcactg aatgccaaagg gaacagaagc acccgcgagta  
60 541 gtgacagagg aggaggatga tgatgaagag actgctcctc ccgttattgc cccgcgaccg  
601 gatcatacga aatcaattta cacacgggtc gtaattgacc ctgttcctgc accagtgggt  
661 gattcacatg ttgatggtgc tgccaagtct ttagacaaac agaaaaagaa gcctaagatg  
721 acagatgaag agattatgga gaaattaaga actatcgtga gcataggtga ccctaagaaa



```

5  781 aaatatacaa gatatgaaaa aattggacaa ggggcttctg gtacagtttt cactgctact
    841 gacgttgacac tgggacagga ggttgctatc aaacaaatta atttacagaa acagccaaag
    901 aaggaactga tcattaacga gattctgggtg atgaaagaat tgaaaaatcc caacatcggt
    961 aacttttttg acagttacct ggtaggagat gaattgtttg tggatcatgga ataccttgct
10 1021 ggggggtcac tcaactgatgt ggtaacagaa acagcttgca tggatgaagc acagattgct
    1081 gctgtatgca gagagtgttt acaggcattg gagttttttac atgctaataca agtgcaccac
    1141 agagacatca aaagtgacaa tgtactttttg ggaatggaag gatctgttaa gctcactgac
    1201 tttgggtttct gtgcccagat cacccttgag cagagcaaac gcagtaccat ggtcggaacg
    1261 ccatactgga tggcaccaga ggtgggttaca cggaaagctt atggccctaa agtcgacata
15 1321 tgggtctctgg gtatcatggc tattgagatg gtagaaggag agcctccata cctcaatgaa
    1381 aatcccttga gggccttgta cctaatagca actaatggaa ccccaagaact tcagaatcca
    1441 gagaaacttt ccccaatatt tcgggatttc ttaaactgat gtttggaagt ggatgtggaa
    1501 aaaaggggtt cagccaaaga attattacag catcctttcc tgaactggc caaaccgtta
    1561 tctagcttga caccactgat catggcagct aaagaagcaa tgaagagtaa ccgttaacat
20 1621 cactgctgtg ggctcactat cttttttcca ttttctacaa gaagcctttt agtatatgaa
    1681 aatgatgact ctggtggggg ttttaagaaa tgggtctgcat aacctgaatg aaagaaggaa
    1741 atgactattc tctgaagaca accaagagaa aattggaaaa gacaaggtat gactttgtta
    1801 tgaacccttg ctttttagggg tccaggaagg gatttgtggg acttgaattc actaggctta
    1861 ggtctttcag gaaacaggct atcaggggca tttatcatgt gtgagattgg attctacttg
25 1921 ggtgatttgg tggatagacc catgaatggc ccctgggggt tttcaatctt ggattggagg
    1981 tgggggtttc agagtgttgc cactctagc tcctctccc
```

25

DEFINITION Human cyclin-dependent kinase 4 (CDK4) gene,  
complete cds.

ACCESSION U37022

VERSION U37022.1 GI:1353415

30 FEATURES Location/Qualifiers

source

1..4233

/organism="Homo sapiens"

/db\_xref="taxon:9606"

/chromosome="12"

35 /map="12q13"

/clone="24C7"

mRNA

join(85..291,729..965,1123..1258,1376..1543,1701..1810,  
2949..2999,3135..3270,3836..4233)

40 /gene="CDK4"

/product="cyclin-dependent kinase 4"

exon

85..291

/gene="CDK4"

gene

85..4233

45 /gene="CDK4"

exon

729..965

/gene="CDK4"

CDS

join(748..965,1123..1258,1376..1543,1701..1810,2949..2999,  
3135..3270,3836..3928)

50 /gene="CDK4"

/codon\_start=1

/product="cyclin-dependent kinase 4"

/protein\_id="AAC50506.1"

55 /db\_xref="GI:1353416"

/translation="MATSRYPVAEIGVGAYGTVYKARDPHSGHFVALKSVRVPNGGG

60 GGGGLP1STVREVALLRRLAEFHPNVVRLMDVCATSRTDREIKVTLVFEHVDQDLRT

YLDKAPPPGLPAETIKDLMRQFLRGLDFLHANCIVHRDLKPENILVTSGGTVKCLADFG

LARIYSYQMALTPVVVTLWYRAPEVLLQSTYATPVDMMWSVGCIFAEMFRRKPLFCGNS

EADQLGKIFDLIGLPPEDDWPRDVSILPRGAFFPRGPRPVQSVVPEMEESGAQLLLEML

TFNPHKRISAFRALQHSYLHKDEGNPE"

	BASE COUNT	945 a	1050 c	1101 g	1124 t	13 others	
5	ORIGIN						
	1	ccctcctccc	agtcgaagca	cctcctgtcc	gcccctcagc	gcatgggtgg	cgggtcacgtg
	61	cccagaacgt	ccggcggttcg	ccccgcctcc	ccagtttccg	cgcgcctctt	tggcagctgg
	121	tcacatggtg	aggggtggggg	tgagggggcc	tctctagctt	gcggcctgtg	tctatggtcg
	181	ggccctctgc	gtccagctgc	tccggaccga	gctcgggtgt	atggggccgt	aggaaccggc
10	241	tccggggccc	cgataacggg	ccgccccac	agcaccgccg	gctggcgtga	ggtaagtga
	301	gtcccttccc	aggaatgaga	accagtcccc	gccccctca	cagctttcca	cgcgttcgtt
	361	tcgcgagctg	gttatggaag	ggtcgctcaa	gggcgggaag	tggggccttt	gtgggtcatgg
	421	gaaagtataa	ttttagggac	tgaggtgtag	gatcttcgat	gcaaggcatg	tgtcatgtgt
	481	gatctttgtg	cggggcgcga	ttgtcccaa	ggaaaaagcg	ttttctattg	cagggcctca
15	541	cgtggctgga	gggggttgga	ttgagtcatt	gtgttatctc	tggggccggc	cccaaggaaag
	601	actgggagcg	ggggatggga	tgctgggtgt	gttctttgct	cttttttttt	gggagtcctt
	661	ttgttgctgc	aggtcatacc	atcctaactc	tgtaagcgac	ttttggtgat	aggagtctgt
	721	gattgtaggg	tctcccttga	tctgagaatg	gctacctctc	gatatgagcc	agtggctgaa
	781	attggtgtcg	gtgcctatgg	gacagtgtac	aaggccccgtg	atccccacag	tggccacttt
20	841	gtggccctca	agagtgtgag	agtccccaat	ggaggaggag	gtggaggagg	ccttcccatc
	901	agcacagttc	gtgaggtggc	tttactgagg	cgactggagg	cttttgagca	tcccaatggt
	961	gtccgggtgag	aaggtgggtg	aggggtgggc	gtggggagta	aagggaaaaag	acagcctata
	1021	ggtgggggtgt	gatgatctgt	agagaagtgg	ggaccctgag	gaaataatga	gaggccatgt
	1081	tgggttaaag	gggattgaaa	agtgagcatt	tactctgggtc	aggctgatgg	acgtctgtgc
25	1141	cacatcccga	actgaccggg	agatcaaggt	aaccctgggtg	tttgagcatg	tagaccagga
	1201	cctaaggaca	tatctggaca	aggcaccctc	accaggcttg	ccagccgaaa	cgatcaaggt
	1261	gagtgggggt	ggtaggcatt	gagaggtgga	ttagggacctt	tgtagttagaa	ccttctggga
	1321	tttcagggtat	ggtgcctagt	ttccagtgca	tctgtacctc	cccctttgaa	actaggatct
	1381	gatgcgccag	tttctaagag	gcctagattt	ccttcatgcc	aattgcatcg	ttcaccgaga
30	1441	tctgaagcca	gagaacattc	tggtgacaag	tggtggaaca	gtcaagctgg	ctgactttgg
	1501	cctggccaga	atctacagct	accagatggc	acttacaccc	gtgggtcagta	gaaagatggt
	1561	acaaaaatgg	gttctggttg	ggaataggag	agtgtattgc	cgtagcaatt	gagaagtcat
	1621	tgcttctcatg	tggttcagtca	agcaagtgtg	gtttcatggt	aaccctgggg	gtccccatcc
	1681	attcttcccta	ttcccttttag	gttggttacac	tctggtaccg	agctcccgaa	gttcttctgc
35	1741	agtcacacata	tgcaacacct	gtggacatgt	ggagtgttgg	ctgtatcttt	gcagagatgt
	1801	ttcgtcgaaa	gtatgggacc	cacataccct	ggactacctt	gaattcccca	aatcgcttgt
	1861	tcataaaacca	catccatacc	ttgcccattc	tttttttttg	agaccagggc	ttgctgtggt
	1921	gcccaggctg	gattgcaatg	gcattgacac	agctcactgc	agcttcaacc	tcctgggctc
	1981	aagtgatcct	cccatctcag	cttcccaact	agctgacact	acaggcacgc	acctccatgc
40	2041	ttggctagtt	tgtaaatatt	tttatagaga	tggggtctca	gtatatgtcc	caggctggtc
	2101	ttgaactctt	gcactcaagc	aatcctccca	cccctacctc	ccaaagtagc	ataagctact
	2161	gcatctggcc	ccattctttt	acttgcgtac	tactaacttg	cccatagcag	aaagctctga
	2221	aatgttctgg	aattaggaac	ttcatatccc	tttattctct	ttatttttta	tttatttatt
	2281	tattttattta	tttatttatt	gagataaagg	ttcactctgn	naccaggct	ggagtncaag
45	2341	ggcccaatta	nagctcactg	tancctctac	ctcctgggct	aaagmaatcc	tcccatctca
	2401	gccccttgag	tanctgagac	taaagggtgca	cgccaccatg	actggctttt	ttttttttta
	2461	gatggagtct	tgctctgtcg	ccaggctgga	gtgcagtagt	gcgatctctg	ctcactgcaa
	2521	cctccacctc	ccagattcaa	gcaattctct	tgactcagcc	tcccaagtag	ctgggaccac
	2581	aggtgcacgc	caccatgctc	agctaatttt	tgtactttta	gtaatgacag	gtttcaccat
50	2641	gttggccagg	atggtctcga	tctcttgacc	tcatgatcca	cccacatcag	actcccaaag
	2701	tgctaggatt	acaggcgtga	gcnnnngcac	ctggcatttc	ttttttttta	aaaaaagaga
	2761	caaggctctg	cttgcccagg	ctgatctaga	actcctgggc	tcaagcagtc	ctctcacctc
	2821	agcatcccaa	agtgtctgaa	ttgttggcct	ttattcccta	tacttccctat	tttgagccac
	2881	taagcagtaa	ccattcaact	aagatatctt	tgaaaatgac	tgctacctta	tatcccttct
55	2941	caccttaggc	ctctcttctg	tggaaaactct	gaagccgacc	agttgggcaa	aatctttgag
	3001	taagtgaaca	acatgggaga	aaaagatttt	ctattctgag	tcctctttct	gctgaaccca
	3061	ggatggcaac	tggtctctgc	atggggatgg	gaactggagg	accctcctga	ccagagtctc
	3121	cctgtccccc	acagcctgat	tgggctgcct	ccagaggatg	actggcctcg	agatgtatcc
	3181	ctgccccgtg	gagcctttcc	ccccagaggg	ccccgcccag	tgcatcggtg	ggtacctgag
60	3241	atggaggagt	cgggagcaca	gctgctgctg	gtaactggag	atggctgtgg	gcacagggaa
	3301	agaaatagag	actggggaaa	gaaatagagc	agtatgcagg	gccctggcca	ctgtggttaa
	3361	tgaactcttg	ttggtagatg	gtctgtagtt	tttattacag	ctgcaaatag	ccaccacag
	3421	agaaggatat	agaagagaac	ccatcctggc	tgggcacggg	ggctcacggc	gttaatccca
	3481	gcacttttggg	aggccaaggt	gggcgtatca	cctgaggtca	ggagttcgag	accagcctgg

12/17

```
3541 ccaacatggt gaaacctcgt ctctactaaa agtacaaaaa taagccgggg gtggtggcac
3601 acgcctgtaa tctcagctac ttgggaggct gagataggag aatcacttca actcaggagg
3661 cggagggttg agtgagctga gttcatacca ttggcactcc agcctgggtg atagagcgag
3721 actccgtctn caaaaaaaaa aaaaaagaaa aaagaagaaa gctcatccca ggtattgttg
5 3781 tgggtggcag aagctgtttt cttcatgggt ttctgacctt tgcctctccc ctcaggaaat
3841 gctgactttt aaccacacac agcgaatctc tgcctttcga gctctgcagc actcttatct
3901 acataaggat gaaggtaatc cggagtggagc aatggagtgg ctgccatgga aggaagaaaa
3961 gctgccattt cccttctgga cactgagagg gcaatctttg cctttatctc tgaggctatg
10 4021 gagggtcctc ctccatcttt ctacagagat tactttgctg ccttaatgac attcccctcc
4081 cacctctcct tttgaggctt ctccttctcc ttcccatttc tctacactaa ggggtatggt
4141 ccctcttgct cctttcccta cctttatatt tggggtcctt ttttatacag gaaaaacaaa
4201 accaaaagaa awaatggccc tttttttttt ttt
```

15

DEFINITION Human MEK5 mRNA, complete cds.

ACCESSION U25265

VERSION U25265.1 GI:1255719

20 FEATURES

Location/Qualifiers

source

1..2083

/organism="Homo sapiens"

/db\_xref="taxon:9606"

/clone="3-2"

25

/tissue\_type="brain"

/clone\_lib="cDNA Lambda ZAP (Invitrogen)"

/dev\_stage="fetal"

CDS

297..1613

/codon\_start=1

30

/product="MEK5"

/protein\_id="AAA96146.1"

/db\_xref="GI:1255720"

/translation="MLWLALGPFPMENQVLVIRIKIPNSGAVDWTVHSGPQLLFRDV

35

LDVIGQVLPEATTTA FEYEDGDGRITVRSDEEMKAMLSYYYSTVMEQQVNGQLIEPL

QIFPRACKPPGERNIHGLKVNTRAGPSQHSSPAVSDSLPSNSLKKSSAELKKILANGQ

40

MNEQDIRYRDTLGHGNGGT VYKAYHVPSGKILAVKVILLDTLELQKQIMSELEILYK

CDSSYIIGFYGAFFVENRISICTEFMDGGS LDVYRKMP EHV LGRIAVAVVKGLTYLWS

45

LKILHRDVKPSNMLVNTRGQVKLCDFGVSTQLVNSIAKTYVGTNAYMAPERISGEQYQ

IHSDVWSLGISFMEIQKNQGS LMPLQLLQCIVDEDS PVLPVGEFSEPFVHFITQCMRK

QPKERPAP EELMGHPFVQFNDGNA AVVSMWVCRAL EERRSQQGPP"

BASE COUNT

518 a

532 c

512 g

521 t

ORIGIN

50

1 agcgttcgct caactccaga accttccgac ctccgctagt tcttgccggc ctttgcgccg

61 ttcccgggtgc accctccccg ggagacacct cagacccccg acagcctggg caggctcggt

121 gcctgcgggt gcgttcctga tcacccctcc cctcttccct cccctcatc ctccattccc

181 ttgttttcac cctctgtcct ctgcccgta ctccccttgt cacctcttgg agccccctcc

241 taaccagcgg ccagtgggtt tccataccc caggatgtga gcctctttaa cctgtaatgc

55

301 tgggtcagc cctgggcccc tttcctgcca tggagaacca ggtgctggtg attcgcatca

361 agatcccaaa tagtggcgcg gtggactgga cagtgcactc cgggcgcgag ttactcttca

421 gggatgtgct ggatgtgata ggccagggtc tgcctgaagc aacaactaca gcatttgaat

481 atgaagatga agatggtgat cgaattacag tgagaagtga tgaggaaatg aaggcaatgc

541 tgtcatatta ttattccaca gtaatggaac agcaagtaaa tggacagtta atagagcctc

60

601 tgcagatatt tccaagagcc tgcaagcctc ctggggaacg gaacatacat ggcctgaagg

661 tgaatactcg ggccggaccc tctcaacaca gcagcccagc agtctcagat tcaactccaa

721 gcaatagctt aaagaagtct tctgctgaac tgaaaaaaat actagccaat ggccagatga

781 atgaacaaga catacgatat cgggacactc ttggtcatgg caacggaggc acagtctaca

841 aagcatatca tgtcccagat gggaaaatat tagctgtaaa ggtcatacta ctagatatta

```

    901 cactggaact tcagaagcaa attatgtctg aattggaaat tctttataag tgcgattcat
    961 catatatcat tggattttat ggagcatttt ttgtagaaaa caggatttca atatgtacag
1021 aattcatgga tgggggatct ttggatgtat ataggaaaaat gccagaacat gtccttgga
1081 gaattgcagt agcagttgtt aaaggcctta cttatttgtg gagtttaaag attttacata
  5 1141 gagacgtgaa gccctccaat atgctagtaa acacaagagg acagggttaag ctgtgtgatt
    1201 ttggagttag cactcagctg gtgaattcta tagccaagac gtatgttgga acaaatgctt
    1261 atatggcgcc tgaaaggatt tcaggggagc agtatggaat tcattctgat gtctggagct
    1321 taggaatctc ttttatggag attcagaaaa accagggatc tttaatgcct ctccagcttc
    1381 tgcagtgcac tgttgatgag gattcgcccg tccttcagct tggagagttc tcggagccat
  10 1441 ttgtacattt catcactcag tgtatgcgaa aacagccaaa agaaaggcca gcacctgaag
    1501 aattgatggg ccacccgttc atcgtgcagt tcaatgatgg aaatgccgcc gtggtgtcca
    1561 tgtgggtgtg ccgggcgctg gaggagaggc ggagccagca gggggccccc tgaggctgcc
    1621 gcagggcact gaaagcccag gaccagtaac caaggagAAC aaccaccccg tcgcccttct
    1681 ccgtatgctg cctgcgccag aagagctttg ctgggccctg gcttcctctg cctcgccctc
  15 1741 acctctgtca gcaggtggcc ttgcctgggg agcccatgt gtggcccacc ccaccaggcc
    1801 atccccatac cttctggttt gaaggcgctg acactggcag agaggtaaag ggtggggcat
    1861 tgagaatgga ggctcccagg gtccctgccc acttctgttt tcctaagtgt tttctctata
    1921 aagggtcagg cccgtcagca tactgatgg gaataaaagt attaatgctt tgtgacagcc
    1981 tctgcctgaa aactggacag aaggaccag aggtgttctt tcattttctc tcttacctcc
  20 2041 aatctttccc ctttcaagct acaggtaaag gctctaccac cat
```

25 DEFINITION Homo sapiens a disintegrin and metalloproteinase domain 22

(ADAM22), transcript variant 3, mRNA.

ACCESSION NM\_016351

VERSION NM\_016351.2 GI:11497041

30 FEATURES Location/Qualifiers

source

1..3314

/organism="Homo sapiens"

/db\_xref="taxon:9606"

/chromosome="7"

/map="7q21"

35

gene

1..3314

/gene="ADAM22"

/note="MDC2"

/db\_xref="LocusID:53616"

40

CDS

/db\_xref="MIM:603709"

47..2659

/gene="ADAM22"

/note="isoform 3 is encoded by variant 3"

/codon\_start=1

45

/db\_xref="LocusID:53616"

/db\_xref="MIM:603709"

/product="a disintegrin and metalloproteinase domain

22,

isoform 3 preproprotein"

50

/protein\_id="NP\_057435.1"

/db\_xref="GI:7706613"

/translation="MQAAVAVSVPFLLLCVLGTCPPARCGQAGDASLMELEKRKENRF

55 VERQSIVPLRLIYRSGGEDES RHDALDTRVRGDLGGRQLTHVDQASFQVDAFGTSFIL

DVVLNHDLLSSEYIERHIEHGGKTVEVKGGEHCYYQGHIRGNPDSFVALSTCHGLHGM

FYDGNHTYLIEPEENDTTQEDFHFHSVYKSRLFEFSLDDLPSFQQVNITPSKFILKP

60

RPKRSKRQLRRYP RNVEETKYIELMIVNDHLMFKKRLSVVHTNTYAKSVVNMADLI

YKDQLKTRIVLVAMETWATDNKFAISENPLITLREFMKYRRDFIKEKSDAVHLFSGSQ

FESSRSGAAYIGGICSLKGGGVNEFGKTDLMVTLAQSLAHNIGIISDKRKLASGEC

KCEDTWSGCIMGDTGYLPPKFTQCNIIEYHDFLNSGGGACLFNKPSKLLDPPECNGN

FIETGEECDCGTPAECVLEGAECCKKCTLTQDSQCS DGLCKKCKFQPMGTVCRAVN

DCDIRETCSGNSSQCAPNIHKMDGYSCDGVQIGCFGGRCKTRDRQCKYIWGQKVTASD

KYCYEKLNIEGTEKGNCGKDKDTWIQCNRDVLGCGYLLCTNIGNIPRLGELDGEITST

LVVQQGRTLNCSSGGHVKLEEDVDLGYVEDGTPCGPQMMCLEHRCPLVASFNFSTCLSS

KEGTICSGNGVCSNELKVCNHRWIGSDCNTYFPHNDDAKTGITLSGNGVAGTNIIIG

IIAGTILVLALILGITAWGYKNYREQRSNGLSHSWSERIPDTKHISDICENGRPRSNS

WQGNLGGNKKKIRGKRFRPRSNSTETLSPAKSPSSSTGSIASSRKYPYPMPLPDEDK

KVNQRSARLWETSI"

BASE COUNT 1000 a 641 c 779 g 891 t 3 others

ORIGIN

1 catgaggagc tgagcgtctc gggcgaggcg ggctgacggc agcaccatgc aggcggcagt

61 ggctgtgtcc gtgcccttct tgctgtctct tgtcctgggg acctgccctc cggcgcgctg

121 cggccaggca ggagacgcct cattgatgga gctagagaag aggaaggaaa accgcttcgt

181 ggagcgccag agcatcgtgc cactgcgcct catctaccgc tcgggcggcg aagacgaaag

241 tcggcacgac gcgctcgaca cgcgggtgcg gggcgacctc ggtggccggc agttgactca

301 tgttgaccaa gcaagcttcc aggttgatgc ctttggaacg tcattcattc tcgatgtcgt

361 gctaaatcat gatttgctgt cctctgaata catagagaga cacattgaac atggaggcaa

421 gactgtggaa gttaaaggag gagagcactg ttactaccag ggccatatcc gaggaaaccc

481 tgactcattt gtgcattgtt caacatgcca cggacttcat gggatgttct atgacgggaa

541 ccacacatat ctcattgagc cagaagaaaa tgacactact caagaggatt tccattttca

601 ttcagtttac aaatccagac tgtttgaatt ttccttggat gatcttccat ctgaatttca

661 gcaagtaaac attactccat caaaatttat tttgaagcca agacaaaaaa ggagtaaacg

721 gcagcttcgt cgatatcctc gtaatgtaga agaagaaacc aaatacattg aactgatgat

781 tgtgaatgat caccttatgt ttaaaaaaca tcggctttcc gttgtacata ccaataccta

841 tgcgaaatct gtggtgaaca tggcagattt aatatataaa gaccaactta agaccaggat

901 agtattgggt gctatggaaa cctgggcgac tgacaacaag tttgccatat ctgaaaatcc

961 attgatcacc ctacgtgagt ttatgaaata caggagggat tttatcaaag agaaaagtga

1021 tgcagttcac cttttttcgg gaagtcaatt tgagagtagc cggagcgggg cagcttatat

1081 tggtgggatt tgctcggtgc tgaaggagg aggcgtgaat gaatttggga aaactgattt

1141 aatggctgtt acacttgccc agtcattagc ccataatatt ggtattatct cagacaaaag

1201 aaagttagca agtgggtgaat gtaaatgcga ggacacgtgg tcgggtgca taatgggaga

1261 cactggctat tatcttccta aaaagttcac ccagtgtaat attgaagagt atcatgactt

1321 cctgaatagt ggaggtgggt cctgcctttt caacaaacct tctaagcttc ttgatcctcc

1381 tgagtgtggc aatggcttca ttgaaactgg agaggagtgt gattgtggaa ccccggccga

1441 atgtgtcctt gaaggagcag agtgtgtgaa gaaatgcacc ttgactcaag acctctaatg

1501 cagtgcagggt ctttgctgta aaaagtgcga gtttcagcct atgggcactg tgtgccgaga

1561 agcagtaaat gattgtgata ttcgtgaaac gtgctcagga aattcaagcc agtgtgcccc

1621 taatattcat aaaatggatg gatattcatg tgatgggtgt caggggaattt gctttggagg

1681 aagatgcaaa accagagata gacaatgcaa atacatttgg gggcaaaaag tgacagcatc

1741 agacaaatat tgctatgaga aactgaatat tgaaggagc gagaagggta actgtgggaa

1801 agacaaagac acatggatac agtgaacaaa acgggatgtg ctttgtggtt accttttgtg

1861 taccaatatt ggcaatatcc caaggcttgg agaactcgat ggtgaaatca catctacttt

1921 agttgtgcag caaggaagaa cattaaactg cagtgggtgg catgttaagc ttgaagaaga

1981 tgtagatctt ggctatgtgg aagatgggac accttgtggt ccccaaatga tgtgcttaga

2041 acacaggtgt cttcctgtgg cttctttcaa ctttagtact tgcttgagca gtaaagaagg

2101 cactatttgc tcaggaaatg gagtttgacg taatgagctg aagtgtgtgt gtaacagaca

2161 ctggataggt tctgattgca acacttactt ccctcacaat gatgatgcga agactgggat

2221 cactctgtct ggcaatgggt ttgctggcac caatatcata ataggcataa ttgctggcac

2281 catttttagtg ctggccctca tattaggaat aactgcgtgg ggttataaaa actatcgaga

2341 acagaggtca aatgggctct ctcattcttg gagtgaaggg attccagaca caaaacatat

2401 ttcagacatc tgtgaaaatg ggcgacctcg aagtaactct tggcaaggta acctggggag

2461 caacaaaaag aaaatcagag gcaaaagatt tagacctcgg tctaattcaa ctgagacttt

2521 atctcctgcc aaqtctcctt cttcatcaac tgggtctatt gcctccagca gaaaataccc

```
2581 ttacccaatg cctccacttc ctgatgagga caagaaagtg aaccgacaaa gtgccaggct
2641 atggggagaca tccattttaag atcaactggt tacatgtgat acatcgaaaa ctgtttactt
2701 caacttttat agaaacccag gctcatggaa tcatcgaaa tctatctgct cttcagacaa
2761 tacgaagacc ctctgagatg ctacagagga gaggaagcgg agtttcacat ctggttacca
5 2821 ttttcttttt gtcataggct taggatttaa ctaaccatga aaagaactac tgaaatatta
2881 cactataaca tggaacaata aaggtagctg tatgttaatg gataatccgc atgacagata
2941 atatgtagaa atattcataa agttaactca catgacccaa atgtagcaag tttcctaagg
3001 tacaatagtg gattcagaac ttgacgttct gaggcacatc ctcactgtaa acagtaatgc
3061 tatatgcatg aagcttctgt ttattgtttt ccatatttaa ggaaacaaca tcccataata
10 3121 gaaatgagca tgcagggcta aggcataatg gatttttctg caggacttta aagctttgaa
3181 aggccaatat cccataggct aactttaaac atgtattttt atttttgntt tgntttttac
3241 ttttcatatt tatattagca tacaaggaca attgnatata tgtaacattt ttaaaatttt
3301 aaaaaaaaaa aaaa
```

15

```
DEFINITION Homo sapiens a disintegrin and metalloproteinase domain 17
(tumor necrosis factor, alpha, converting enzyme) (ADAM17),
20 transcript
variant 1, mRNA.
ACCESSION NM_003183
VERSION NM_003183.3 GI:11497003
25 FEATURES
Location/Qualifiers
source 1..3523
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="2"
30 /map="2p25"
gene 1..3523
/gene="ADAM17"
/note="TACE; CSVP; CD156b"
/db_xref="LocusID:6868"
35 /db_xref="MIM:603639"
CDS 165..2639
/gene="ADAM17"
/note="isoform 1 is encoded by transcript variant 1;
TNF-alpha converting enzyme; snake venom-like
40 protease"
/codon_start=1
/db_xref="LocusID:6868"
/db_xref="MIM:603639"
/product="a disintegrin and metalloproteinase domain
45 17,
isoform 1 preproprotein"
/protein_id="NP_003174.2"
/db_xref="GI:11497004"
50 /translation="MRQSLFLTSVVPFVLAPRPDDPGFGPHQRLEKLDLLSDYDI
LSLSNIQQHSVRKRDLOTSTHVETLLTFSALKRHFKLYLTSSTERFSQNFKVVVVDGK
NESEYTAKWQDFFTGHVVGEPDSRVLAHIRDDDVIIIRINTDGAEYNIEPLWRFVNDTK
55 DKRMLVYKSEDIKNVSRLQSPKVCGYLKVDNEELLPKGLVDREPPEELVHRVKRRADP
DPMKNTCKLLVADHRFYRYMGRGEESTTTNYLIELIDRVDDIYRNTSWDNAGFKGYG
60 IQIEQIRILKSPQEVKPGKEKHYNMAKSYPNEEKDAWDVKMLLEQFSFDIAEEASKVCL
AHLFTYQDFDMGTLGLAYVGSPRANSHGGVC PKAYYSPVGKKNIYLN SGLTSTKKNY GK
TILTKEADLVTTHELGHNF GAEHDPDGLAE CAPNEDQGGKYVMYPIAVSGDHENNKMF
```

SNCSKQSIYKTIESKAQECFQERSNKVCGNSRVDEGEECDPGIMYLNNDTCCNSDCTL

KEGVQCSDRNSPCKKNCQFETAQKKCQEAINATCKGVSYCTGNSSECPPPGNAENDTV

5

CLDLGKCKDGGKCIFFCEREQQLSCACNETDNSCKVCCRDLSGRVCVPYVDAEQKNLFL

RKGKPCTVGFCDMNGKCEKRVQDVIERFWDFIDQLSINTFGKFLADNIVGSVLVFSLI

10

FWIPFSILVHCVDKKLDKQYESLSLFHPSNVEMLSMDSASVRIKPPAPQTPGRLQ

PAPVIPSAPAAPKLDHQRMDTIQEDPSTDHMDDEGFEKDPFPNSSTAAKSFEDLTDH

PVARSEKAASFKLQRQNRVNSKETEC"

BASE COUNT 1025 a 685 c 845 g 968 t

15

ORIGIN

1	tcgagcctgg	cggtagaatc	ttcccagtag	gcggcgcggg	agggaaaaga	ggattgaggg
61	gctaggccgg	gcggtatccg	tcctccccc	atgtgagcag	ttttccgaaa	ccccgtcagg
121	cgaaggctgc	ccagagaggt	ggagtcggtg	gcggggccgg	gaacatgagg	cagtctctcc
181	tattcctgac	cagcgtgggt	cctttcgtgc	tggcgccggc	acctccggat	gacccgggct
241	tcggccccc	ccagagactc	gagaagcttg	attccttgct	ctcagactac	gatattctct
301	ctttatctaa	tatccagcag	cattcggtta	gaaaaagaga	tctacagact	tcaacacatg
361	tagaaacact	actaaccttt	tcagctttga	aaaggcattt	ttaattatac	ctgacatcaa
421	gtactgaacg	tttttcacaa	aatttcaagg	tcgtgggtgg	ggatggtaaa	aacgaaagcg
481	agtacactgc	aaaatggcag	gacttcttca	ctggacacgt	ggttgggtgag	cctgactcta
541	gggttctagc	ccacataaga	gatgatgatg	ttataatcag	aatcaacaca	gatggggccg
601	aatataacat	agagccactt	tggagatttg	ttaatgatac	caaagacaaa	agaatgttag
661	tttataaatc	tgaagatatc	aagaatgttt	cacgtttgca	gtctccaaaa	gtgtgtgggt
721	atttaaaagt	ggataatgaa	gagttgctcc	caaaagggtt	agtagacaga	gaaccacctg
781	aagagcttgt	tcacgagtg	aaaagaagag	ctgaccacga	tcccatgaag	aacacgtgta
841	aattattggg	ggtagcagat	catcgcttct	acagatacat	gggcagaggg	gaagagagta
901	caactacaaa	ttacttaata	gagctaattg	acagagttga	tgacatctat	cggaaacactt
961	catgggataa	tgcaggtttt	aaaggctatg	gaatacacat	agagcagatt	cgcattctca
1021	agtctccaca	agaggtaaaa	cctggtgaaa	agcactacaa	catggcaaaa	catgtaccac
1081	atgtacaaaa	ggatgctttg	gatgtgaaag	tgttgctaga	gcaatttagc	tttgatatag
1141	ctgaggaagc	atctaaagtt	tgcttggcac	accttttcac	ataccaagat	tttgatatgg
1201	gaactcttgg	attagcttat	gttggctctc	ccagagcaaa	cagccatgga	gggtgttgct
1261	caaaggctta	ttatagccca	gttgggaaga	aaaatatcta	tttgaatagt	ggtttgacga
1321	gcacaaagaa	ttatggtaaa	accatcctta	caaaggaagc	tgacctggtt	acaactcatg
1381	aattgggaca	taattttgga	gcagaacatc	atccggatgg	tctgcagaaa	ttgtgcccga
1441	atgaggacca	gggagggaaa	tatgtcatgt	atcccatagc	tgtgagtggc	gatcacgaga
1501	acaataagat	gttttcaaac	tgcagtaaac	aatcaatcta	taagaccatt	gaaagtaagg
1561	cccaggagtg	ttttcaagaa	cgcagcaata	aagtttgtgg	gaactcgagg	gtggatgaag
1621	gagaagagtg	tgatcctggc	atcatgtatc	tgaacaacga	cacctgctgc	aacagcgact
1681	gcacgttgaa	ggaaggtgtc	cagtgacgtg	acaggaacag	tccttgctgt	aaaaactgtc
1741	agttttagac	tgcccagaag	aagtgcagg	agggcattaa	tgctacttgc	aaaggcgtgt
1801	ccttctgcac	aggtaatagc	agtgatggcc	cgccctccagg	aaatgctgaa	aatgacactg
1861	tttgcttgga	tcttggcaag	tgtaaggatg	ggaaatgcat	ccctttctgc	gagagggaa
1921	agcagctgga	gtcctgtgca	tgtaaatgaa	ctgacaactc	ctgcaagggtg	tgctgcaggg
1981	acctttctgg	ccgctgtgtg	ccctatgtcg	atgctgaaca	aaagaactta	tttttgagga
2041	aaggaaagcc	ctgtacagta	ggattttgtg	acatgaatgg	caaatgtgag	aaacgagtac
2101	aggatgtaat	tgaacgattt	tgggatttca	ttgaccagct	gagcatcaat	acttttggaa
2161	agtttttagc	agacaacatc	gttgggtctg	tcctgggttt	ctccttgata	ttttggattc
2221	ctttcagcat	tcttgtccat	tgtgtggata	agaaattgga	ttaacagtat	gaatctctgt
2281	ctctgtttca	ccccagtaac	gtcgaaatgc	tgagcagcat	ggattctgca	tcggttcgca
2341	ttatcaaacc	ctttcctgcy	ccccagactc	caggccgcct	gcagcctgcc	cctgtgatcc
2401	cttcggcgcc	agcagctcca	aaactggacc	accagagaat	ggacaccatc	caggaagacc
2461	ccagcacaga	ctcccatatg	gacgaggtat	ggtttgagaa	ggacccttcc	ccaaatagca
2521	gcacagctgc	caagtcattt	gaggactctc	cggaccatcc	ggtcgccaga	agtgaagagg
2581	ctgcctcctt	taaactgcag	cgtcagaatc	gtgttaacag	caaagaaaca	gagtgcta
2641	ttagttctca	gctcttctga	cttaagtgtg	caaaatattt	ttatagattt	gacctacaaa
2701	tcaatcacag	cttgatattt	gtgaagactg	ggaagtgact	tagcagatgc	tggtcatgtg
2761	tttgaacttc	ctgcaggtaa	acagttcttg	tgtgggttgg	cccttctcct	tttgaagagg
2821	taaggtgaaa	gtgaatctac	ttattttgag	gctttcaggt	tttagttttt	aaaatatctt
2881	ttgacctgtg	gtgcaaaaag	agaaaataca	gctggattgg	gttatgaata	tttacgtttt

2941 tgtaaattaa tctttttatat tgataacagc actgactagg gaaatgatca gttttttttt  
3001 atacactgta atgaaccgct gaatatgaag catttgcat ttatttgtga gaaaagtggga  
3061 atagtttttt tttttttttt ttttttttgc cttcaactaa aaacaaagga gataaattta  
3121 gtatacattg tatctaaatt gtgggtctat ttctagttag taccagagt ttttatgtag  
5 3181 cagggaaaat atatatctaa atttagaaat catttggtt aatatggctc ttcataattc  
3241 taagactaat gctcagaacc taaccactac cttacagtga gggctataca tggtagccag  
3301 ttgaatttat ggaatctacc aactgtttag ggccctgatt tgctgggcag tttttctgta  
3361 ttttataagt atcttcatgt atccctgtta ctgataggga tacatgtctt agaaaattca  
3421 ctattggctg ggagtgggtg ctcatgcctg taatcccagc acttggagag gctgaggttg  
10 3481 cgccactaca ctccagcctg ggtgacagag tgagatctgc ctc

Accession number: NM\_007026

Definition: DUSP14, MKP-L

15 GTGGCCGCGCAGGAGGACGGAGCCCTAACCGCAACCCGCGCCGCGCCGCGCGATTGATTTGTATCCAC  
TGTCACCAGCACTGCTCACTTAGGACTTTCTGGATCCAGACCCAGGCAGCGCACACTGGACTCTTGAGGA  
AGAAGGAGACTCTAATTTTGGATTCTTGGTGGAGGAAAAATAAACACTCTGGTCTTGCCGCCAACGATG  
CAAGTGTGACTGCTGGCGTCTTCATGAGCTCCAGAGGTCACAGCACGCTACCAAGGACTCTCATGGCCCC  
TCGGATGATTTCCGAGGGAGACATAGGAGGCATTGCTCAAATCACCTCCTCTCTATTCTTGGGCAGAGGC  
20 AGTGTGGCCTCCAATCGGCACCTCCTCCAGGCTCGTGGCATCACCTGCATTGTTAATGCTACCATTGAGA  
TCCCTAATTTCAACTGGCCCCAATTTGAGTATGTTAAAGTGCCTCTGGCTGACATGCCGCATGCCCCCAT  
TGGACTGTACTTTGACACCGTGGCTGACAAGATCCACAGTGTGAGCAGGAAGCACGGGGCCACCTTGGTG  
CACTGTGCTGCAGGGGTGAGCCGCTCAGCCACGCTGTGTATCGCGTACCTGATGAAATTCACAACGTGT  
GCCTGCTGGAGGCATACAACTGGGTGAAAGCCCGGCGACCTGTCTATCAGGCCAACGTAGGCTTCTGGAG  
25 GCAACTGATAGACTACGAGCGCCAGCTCTTTGGGAAGTCGACAGTTAAAATGGTACAGACACCTTATGGC  
ATAGTTCCTGACGCTCTATGAGAAGGAGTCCCGACACCTGATGCCCTTACTGGGGGATTTAGTGCCACTGAA  
GCCTGCGTCTCAGCAGCCCGAGCGGGGCCGGCATCTGCTCCCGCCGCTCTGCTCCCTCTCCACTCTCTCTC  
AAATGGCTGACTTCTGGTTCTCCCTCAAGTGTTTTTTACACTGGGTGTTCAAATTTATTTTAAGAGATAG  
GGAGGGAGGGGACATAAAGGGAATGCATACATTGCTAGTCACATTTTTTAAATTAACATTTTGGAATAGT  
30 GTTTATGGAAATCTTTAGCTTTTAAATCATTTTTACCAATTTGAACAGTTTAATAAACTGGTTCTGCTCTC  
TTCTGAATCTCATGCCCTTTGGCACCTTGGTAGGTGCAGGAGGAGCTCAGTGCAAAAATCACTTTGGGGCC  
TCATTAACCTTTAGAGACAAGCTTTGCCCCAGGCTGCGGACCAGACAGATGCTTAGGGAAGGTTGATAC  
CAGCTTCAGTCTCTACTGGATTAGCCCTACTCTTCCCTTTCCCTCCATTATTTAGTGACTCTGTAAGTA  
AGTTAAATACACCTTATTATTTAGCTGTTAAGTAACTATAATGAAATCTGCTGCAAAATCTCTCTTGA  
35 ATCCATGTGCCAGGATTATATTAGCATTATTTTAAATAAATCTATATGCTTAAAAAAAAAAAAAAAAAA  
A



(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number  
**WO 2003/054228 A3**

(51) International Patent Classification<sup>7</sup>: C12Q 1/42, 1/48,  
1/37, 1/68, C07K 16/00, A61K 39/42

(74) Agents: GINZEL, Christian et al.; c/o Zimmermann &  
Partner, Postfach 330 920, 80069 Munich (DE).

(21) International Application Number:  
PCT/EP2002/014578

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZM, ZW.

(22) International Filing Date:  
19 December 2002 (19.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/341,757 21 December 2001 (21.12.2001) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): AXXIMA  
PHARMACEUTICALS AG [DE/DE]; Max-Leb-  
sche-Platz 32, 81377 München (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): SALASSIDIS, Kon-  
stadinos [GR/DE]; Hauptstrasse 14c, 85386 Eching (DE).  
SCHUBART, Daniel [DE/DE]; Avenue Charles de Gaulle  
1c, 13469 Berlin (DE). GUTBROD, Heidrun [DE/DE];  
Weberhofstrasse 8, 82166 Graefelfing (DE). MUELLER,  
Stefan [DE/DE]; Thalkirchnerstrasse 184, 81371 Munich  
(DE). KRAETZER, Friedrich [DE/DE]; Geisenbrunner-  
strasse 51, 81475 Munich (DE). OBERT, Sabine [DE/DE];  
Bellinzonastrasse 17, 81475 Munich (DE).

Published:

— with international search report

(88) Date of publication of the international search report:  
15 January 2004

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*



WO 2003/054228 A3

(54) Title: HUMAN CELLULAR ENZYMES AS TARGETS AGAINST HEPATITIS C VIRUS INFECTIONS

(57) Abstract: The present invention relates to human cellular protein kinases, metalloproteases and one phosphatase: beta-adren-  
ergic receptor kinase 1 (NM 001619), Mitogen activated protein kinase activated protein kinase 5 (AF032437), Insulin-stimulated  
protein kinase 1 (U08316), Discoidin domain receptor family, member 1 (NM 013994), Protein Kinase C, mu (X75756), Protein  
Kinase C, theta (L01087), AMP-activated protein kinase beta 2 subunit (AJ 224538), JNK2 (U09759), Human p21-activated protein  
kinase 2 (U24153), cyclin-dependent kinase 4 (U37022), MEK5 (U25265), MKP-L (NM 007026), ADAM22 (NM 016351), and  
ADAM17 (U92649) as potential targets for medical intervention against Hepatitis C virus (HCV) infections. The present invention  
relates also to a method for the detection of compounds useful for prophylaxis and/or treatment of Hepatitis C virus infections, a  
method for detecting Hepatitis C virus infections in an individual or in cells. Mono- or polyclonal antibodies are disclosed effective  
for the treatment of HCV infections together with methods for treating Hepatitis C virus infections or for the regulation of Hepatitis  
C virus production and/or replication wherein said antibodies may be used. Finally the present invention relates to a solid support  
useful for detecting Hepatitis C virus infections or for screening compounds useful for prophylaxis and/or treatment of HCV infec-  
tions.

# INTERNATIONAL SEARCH REPORT

Additional Application No  
PCT/EP 02/14578

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/42 C12Q1/48 C12Q1/37 C12Q1/68 C07K16/00  
A61K39/42

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12Q C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, EMBASE, MEDLINE, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BRANCH ANDREA D ET AL: "The coming impact of gene expression profiling on the diagnosis and treatment of HCV-associated liver disease." ANTIVIRAL RESEARCH, vol. 52, no. 2, November 2001 (2001-11), pages 173-179, XP002249604 ISSN: 0166-3542 the whole document	1-14
X	BIGGER CATHERINE B ET AL: "DNA microarray analysis of chimpanzee liver during acute resolving hepatitis C virus infection." JOURNAL OF VIROLOGY, vol. 75, no. 15, August 2001 (2001-08), pages 7059-7066, XP002249605 ISSN: 0022-538X the whole document	1-14

-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

30 July 2003

Date of mailing of the international search report

21/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Niemann, F

# INTERNATIONAL SEARCH REPORT

International Application No <b>PCT/EP 02/14578</b>
--

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>YAMASHITA TARO ET AL: "Serial analysis of gene expression in chronic hepatitis C and hepatocellular carcinoma."            BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS,            vol. 282, no. 2,            30 March 2001 (2001-03-30), pages 647-654,            XP002249606            ISSN: 0006-291X            the whole document</p> <p style="text-align: center;">---</p>	1-14
X	<p>HUANG YING ET AL: "A human hepatoma cell line expressing hepatitis C virus nonstructural proteins tightly regulated by tetracycline."            BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS,            vol. 281, no. 3,            2 March 2001 (2001-03-02), pages 732-740,            XP002249607            ISSN: 0006-291X            the whole document</p> <p style="text-align: center;">-----</p>	1-14

# INTERNATIONAL SEARCH REPORT

...ternational application No.  
PCT/EP 02/14578

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.1

Although claims 2, 3. are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claims 6-9 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition

## Continuation of Box I.2

Present claims 6-7, 14 relate to an agent defined by reference to a desirable characteristic or property, namely inhibiting at least aprtially the activity of at least one of the enzymes cited in claim 1.

The claims cover all agents having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such agent. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the agent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to antibody which binds to at least one of the enzymes cited in claim 1 (see claim 8).

Present claims 10-11 relate to an oligonucleotide defined by reference to a desirable characteristic or property, namely being capable of detecting activity of at least one of the enzymes cited in claim 1.

The claims cover all oligonucleotides having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such oligonucleotide. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the oligonucleotide by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to oligonucleotides encoding one of the enzymes cited in claim 1 (see claim 12).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

Invention 1 : 1-14 (all partially)

method for detecting compounds useful for the prophylaxis  
and/or treatment of Hepatitis C virus infection, the method  
comprising contacting a test compound with at least  
beta-adrenergic receptor kinase 1  
method for detecting Hepatitis C virus infection

Inventions 2-14 : 1-14 (all partially)

method for detecting compounds useful for the prophylaxis  
and/or treatment of Hepatitis C virus infection, the method  
comprising contacting a test compound with at least one of  
the enzymes mentioned in claim 1  
method for detecting Hepatitis C virus infection

**THIS PAGE LEFT BLANK**